

Published in final edited form as:

*Int J Obes (Lond)*. 2011 May ; 35(5): 666–675. doi:10.1038/ijo.2010.187.

## Association between the *APOA2* promoter polymorphism and body-weight in Mediterranean and Asian populations. Replication of a gene-saturated fat interaction

Dolores Corella<sup>1,2,3</sup>, E Shyong Tai<sup>4</sup>, Jose V Sorlí<sup>2,3</sup>, Suok Kai Chew<sup>5</sup>, Oscar Coltell<sup>1,6</sup>, Mercedes Sotos-Prieto<sup>2,3</sup>, Antonio García-Rios<sup>3,7</sup>, Ramón Estruch<sup>3,8</sup>, and Jose M. Ordovas<sup>1,3,9,\*</sup>

<sup>1</sup>Nutrition and Genomics Laboratory, JM-USDA Human Nutrition Research Center on Aging at Tufts University, Boston, MA

<sup>2</sup>Genetic and Molecular Epidemiology Unit School of Medicine. University of Valencia, Valencia, Spain

<sup>3</sup>CIBER Fisiopatología de la Obesidad y Nutrición, Instituto de Salud Carlos III, Madrid, Spain

<sup>4</sup>Department of Endocrinology, Singapore General Hospital, Singapore

<sup>5</sup>Epidemiology and Disease Control Division, Ministry of Health Singapore, Singapore

<sup>6</sup>Department of Computing Languages and Systems. Universitat Jaume I. Castellon, Spain.

<sup>7</sup>Reina Sofia University Hospital, Lipids and Atherosclerosis Research Unit, University of Cordoba, Cordoba, Spain

<sup>8</sup>Department of Internal Medicine. Hospital Clinic, Barcelona, Spain

<sup>9</sup>Department of Cardiovascular Epidemiology and Population Genetics, Centro Nacional de Investigaciones Cardiovasculares (CNIC), Madrid, Spain

### Abstract

**Objective**—The *APOA2* gene has been associated with obesity and insulin resistance (IR) in animal and human studies with controversial results. We have reported an *APOA2*-saturated fat interaction determining body mass index (BMI) and obesity in American populations. This work aims to extend our findings to European and Asian populations.

**Methods**—Cross-sectional study in 4602 subjects from 2 independent populations: A high cardiovascular risk Mediterranean population (n=907 men and women; aged 67+/-6 years) and a multiethnic Asian population (n=2506 Chinese, n=605 Malays and n=494 Asian Indians; aged 39+/-12 years), participating in a Singapore National Health Survey. Anthropometric, clinical, biochemical, lifestyle and dietary variables were determined. Homeostasis model assessment of IR (HOMA-IR) was used in Asians. We analyzed gene-diet interactions between the *APOA2* -265T>C polymorphism and saturated fat intake (<or>=22 g/d) on anthropometric measures and IR.

Users may view, print, copy, download and text and data- mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: [http://www.nature.com/authors/editorial\\_policies/license.html#terms](http://www.nature.com/authors/editorial_policies/license.html#terms)

\***Correspondence to:** Jose M. Ordovas. Director, Nutrition and Genomics Laboratory, JM-USDA Human Nutrition, Research Center on Aging at Tufts University, 711 Washington Street, Boston, MA 02111-1524, USA. Tel: (617) 556-3102. Fax: (617) 556-3211, [jose.ordovas@tufts.edu](mailto:jose.ordovas@tufts.edu).

**Financial Disclosures:** None reported.

**Results**—Frequency of CC subjects differed among populations (1%–15%). We confirmed a recessive effect of the *APOA2* polymorphism, and replicated the *APOA2*–saturated fat interaction on body-weight. In Mediterranean individuals, the CC genotype was associated with a 6.8% greater BMI in those consuming a high ( $P=0.018$ ), but not a low ( $P=0.316$ ) saturated fat diet. Likewise, the CC genotype was significantly associated with higher obesity prevalence in Chinese and Asian Indians only with a high-saturated fat intake ( $P=0.036$ ). We also found a significant *APOA2*–saturated fat interaction in determining IR in Chinese and Asian Indians ( $P=0.026$ ).

**Conclusion**—The influence of the *APOA2* –265T>C polymorphism on body-weight-related measures was modulated by saturated fat in Mediterranean and Asian populations.

### Keywords

Obesity; gene-diet interaction; insulin resistance; saturated fat; *APOA2*

## INTRODUCTION

For over two decades, the search for the genetic background underlying common forms of obesity has yielded scores of associations that, for the most part, could not be successfully validated in replication studies (1). Genome-wide association (GWA) studies have led to the discovery of new obesity genes (2–4). Among them, the obesity-associated gene (*FTO*) has now become the first gene with associations thoroughly replicated (2,5,6). Nevertheless, common interactions with environmental factors may have contributed to mask the effect of other single nucleotide polymorphisms (SNPs) on obesity (7). Progress in the investigation and replication of gene-environment interactions for obesity has been slow. However, some steps forward have been accomplished, such as those involving the *FTO* with physical activity (8–10), and the *APOA2* with saturated fat intake (11). In both cases, significant interactions were found and replicated in independent populations. Regarding the *APOA2* locus, several associations with obesity, insulin-resistance (IR) and diabetes have been reported both in animal (12–15) and human studies (16–18). Specifically, the –265T>C promoter polymorphism (rs5082), which has proved to be functional in two independent investigations (19,20), has been significantly associated with obesity-related measures in some human studies (11,19,21–23). In a previous investigation carried out on White Americans participating in the Genetics of Lipid Lowering Drugs and Diet Network (GOLDN) study (21), we observed recessive effects for the –265T>C polymorphism. Homozygous individuals for the –265C allele had higher body mass index (BMI) and obesity risk than did carriers of the T allele. However, we did not find significant associations between this SNP and body-weight measures in the Framingham Study or in the Boston Puerto Rican study, but we did find significant interactions between this SNP and saturated fat intake (11). When saturated fat intake was high (higher than 22 g/d; approx 10% of energy), individuals homozygous for the –265C allele had higher mean BMI than carriers of the T allele (an association which was statistically significant), but no differences in BMI were found when saturated fat intake was low.

Therefore, our objectives were to extend previous findings by: 1) analyzing the association between the *APOA2* –265T>C SNP and body-weight-related variables in a Mediterranean and in a multiethnic Asian population; 2) examining the replication of the previously reported interaction between this SNP and saturated fat intake on body-weight in these populations and 3) exploring the association of this SNP with IR and diabetes and its potential modulation by saturated fat.

## PATIENTS AND METHODS

We analyzed 4,602 subjects from two independent populations. All participants gave their informed consent. The ethics committee of the University of Valencia and the ethics committee of the Singapore General Hospital approved the studies.

### Mediterranean population

We studied 907 unrelated Whites (325 men and 582 women), aged 55–80 years who participated in the PREDIMED (Prevención con Dieta Mediterránea) study and were consecutively recruited in the Valencia Region (Spain) from October 2003 to September 2008. Details of this study have been reported elsewhere (24). Briefly, high-risk participants were selected by physicians in primary care centers. Eligible subjects were elderly community-dwelling persons who fulfilled at least 1 of 2 criteria: type 2 diabetes or 3 or more cardiovascular disease risk factors (current smoking, hypertension, dyslipidemia, overweight, or a family history of premature cardiovascular disease). Here, we included data from the baseline cross-sectional examination.

### Asian population

We studied 3,695 individuals (1,714 males and 1,981 females) who participated in the 1998 Singapore National Health Survey and had genetic data. The detailed methodology of this survey on a nationally representative household sample has been described elsewhere (25). Briefly, the survey protocol was based on the WHO-recommended model for field surveys on diabetes and the WHO MONICA protocol. Individuals representing the Singapore housing population (Chinese, Malays and Indians) aged 18–69 years were selected by disproportionate stratified and systematic sampling. Dietary intake was only assessed in a random sub-sample (1 in 2) of the participants. In the present study, dietary data were available for 1,923 subjects (1,204 Chinese, 412 Malays and 307 Asian Indians). These subjects did not differ in the main variables examined from subjects without dietary data.

### Anthropometric, biochemical, clinical and lifestyle determinations

Anthropometric variables including height, weight and waist were measured in both populations by standard techniques (24,25). Body mass index (BMI) was calculated as weight (kg)/height<sup>2</sup>(m). Obesity was defined as BMI $\geq$ 30kg/m<sup>2</sup> for the Mediterranean population and as BMI $\geq$ 27kg/m<sup>2</sup> for the Asian population (26,27). Fasting triglycerides and HDL-C were measured by standard methods (24,25). In the Asian population, fasting glucose (Boehringer Mannheim, Mannheim, Germany) and fasting insulin (immunoassay using an Abbott AxSYM; Abbott Laboratories, Chicago, IL) were determined in all participants (25). IR (fasting glucose  $\times$  fasting insulin/22.5) was calculated using the homeostasis model assessment of insulin resistance (HOMA-IR) method (28), as a biomarker for IR. In the Asian population, previously diagnosed diabetes was assessed by questionnaire (25). After the fasting sample collection, a 75-g oral glucose tolerance test was taken for all subjects except for previously diagnosed diabetic subjects. After 2 h, glucose (2-h glucose) and insulin (2-h insulin) were measured. Following biochemical determinations, subjects with fasting glucose  $\geq$ 126 mg/dL or 2-h glucose  $\geq$ 200 mg/dL were also classified as having diabetes (29). In the PREDIMED Study, IR was not determined. Diagnosis of type 2 diabetes was based on at least one of the following criteria: i) Current treatment with insulin or oral hypoglycemic drugs; ii) Fasting blood glucose  $\geq$ 126 mg/dl in two measurements; iii) Casual fasting blood glucose  $\geq$  200mg/dl; iv) 2-h glucose  $\geq$ 200 mg/dL (after the oral glucose tolerance test). In the PREDIMED Study, practically all the diabetic subjects (>99%) had previously been diagnosed by the doctor (based on the above criteria) before entering the study. However, in the Asian population an important number of previously unknown diabetes cases (more than 55% of total diabetics) were detected.

In the PREDIMED Study, the baseline examination included assessment of standard cardiovascular risk factors, medication use, socio-demographic factors and lifestyle variables, as previously detailed (24). Physical activity was estimated by the Minnesota Leisure Time Physical Activity questionnaire (30). In Asians, an interviewer-administered questionnaire was used to capture data on sociodemographic factors, smoking, alcohol consumption and physical activity as previously reported (25).

### Dietary intake

In the Mediterranean population food consumption and nutrient intake was determined by a validated food frequency questionnaire (FFQ) (31) and Spanish food composition tables. In Singapore, a validated FFQ was used (32). It comprised 159 individual food items, grouped into 23 main food types and 25 food subtypes to ensure that foods consumed by the three ethnic groups were represented. The food composition database of the Singapore Ministry of Health was used to estimate macronutrient intake.

### Genetic analyses

We performed the *APOA2*-265T>C genotyping (rs5082) using a Taqman assay with allele-specific probes in the ABIPrism 7900HT Sequence Detection System (Applied Biosystems) (21). Quality control measures including positive and negative controls as well as replicated samples (10% at random) were employed. Genotype frequencies were consistent with Hardy-Weinberg equilibrium ( $P=0.10$ ,  $P=0.79$ ,  $P=0.59$ ,  $P=0.20$  for Mediterranean subjects, Chinese, Malays and Asian Indians, respectively).

### Statistical analyses

The Chi square test was used to compare proportions. We applied ANOVA or Student's *t*-test to compare crude means. Multivariate adjustment of continuous variables was carried out by linear regression. Triglycerides and IR were log-transformed. Recessive effects of the *APOA2* SNP were tested and considered in the models. To study gene-diet interactions in determining BMI and body-weight, we used multivariate linear regression models including main effects and interaction terms and control for the same variables that we reported in our previous study (11). We fitted separate models for each population. Saturated fat intake was considered as categorical (low or high). 22g/d was established as the cut-off point based on our previous work (11). We adjusted interaction models for sex, age, ethnicity (when including more than one population), smoking, drinking, diabetes, lipid medication and total energy intake (basic models). The adjustment for lipid medication was not included in the Asian populations because this information was not available. Further adjustments of basic models for physical activity or hypertension were also considered. In addition, adjustment for total fat, MUFA and carbohydrates were carried out when indicated.

Logistic regression models, including main effects and interaction terms, were fitted to test the *APOA2*-SATFAT interaction in determining the odds ratio (OR) of obesity and diabetes. OR and 95% confidence interval (CI) were estimated for each stratum of saturated fat. Multivariate adjustments were undertaken as indicated. Standard regression diagnostic procedures were used to ensure the appropriateness of the fitted models. All reported probability tests were 2-sided. Differences were considered significant at  $P < 0.05$ .

## RESULTS

### Characteristics of the populations

We studied subjects from a Mediterranean and a multiethnic Asian population (Table 1). By design, Mediterranean subjects were older and had a higher prevalence of cardiovascular risk factors. In Singapore, prevalence of diabetes was higher in Asian Indians (19.6%).

Chinese had lower BMI and diabetes. Prevalence of CC subjects was higher in the Mediterranean population (15%). Saturated fat intake was lower in Chinese. We did not find significant associations between the *APOA2* SNP and saturated fat intake either in Chinese or Asian Indians, but a borderline significant association was observed in Malays ( $P=0.08$  Fisher exact test). 100% of the Malay CC subjects ( $n=6$ ) consumed more than 22 g/d of saturated fat vs 59% in T-allele carriers. In the Mediterranean population, we found a significant association ( $P=0.014$ ) in males (84% of the CC males consumed  $\geq 22$  g/d of saturated fat vs 65% in T-allele carriers). No significant association was detected in females.

### Associations between the *APOA2* –265T>C SNP and anthropometric variables

Consistent with our previous findings (11), the effects of this SNP in body-weight were recessive in both Mediterranean subjects and Asian Indians (Figure 1). No associations were found in Chinese or Malays (Table 2). In Asian Indians, homozygous CC subjects had higher means of weight, BMI and waist circumference than T-allele carriers. Prevalence of obesity ( $BMI \geq 27$  Kg/m<sup>2</sup>) was also statistically higher in CC Indian subjects (OR for the CC in comparison with TT+CT subjects was 3.21; 95%CI: 1.45–7.12,  $P=0.004$ ), after multivariate adjustment for sex, age, smoking, drinking and diabetes. Further adjustment for physical activity did not change the estimation ( $P=0.005$ ). In the Mediterranean population, the *APOA2* SNP was significantly associated with weight and BMI after multivariate adjustment for sex, age, smoking, drinking and diabetes. No significant association was found with waist circumference ( $P=0.375$ ). Obesity was not significantly associated with the CC genotype, either unadjusted or after adjustment for sex, age, smoking, drinking and diabetes (OR: 1.28; 95%CI, 0.83–1.96,  $P=0.264$ ).

### Interactions between the *APOA2* –265T>C SNP and saturated fat intake in determining anthropometric variables

According to the interaction model previously reported (11), two categories of saturated fat intake were considered ( $<22$ g/d and  $\geq 22$  g/d). In the Mediterranean population (Figure 2 A), we found a statistically significant interaction between the *APOA2* SNP and saturated fat on BMI in the multivariate adjusted model ( $P$  for interaction: 0.030). Among those within the lower saturated fat strata ( $<22$ g/d), the *APOA2* SNP was not significantly associated with BMI ( $P=0.316$ ). In contrast, the CC genotype was associated with greater BMI ( $P=0.018$ ) in the higher saturated fat strata. Further adjustment for physical activity, total fat intake, MUFA and carbohydrates did not change the statistical significance ( $P: 0.045$ ). In the subgroup analysis by gender, we found that the magnitude of the interaction was higher in women than in men. However, taking into account the lower sample size for men, we cannot confirm differential effects.

In the Asian population, the *APOA2*-saturated fat interactions were not analyzed in Malays because there were no CC subjects that had a saturated fat intake lower than 22g/d. Firstly, the interaction analysis was undertaken in Chinese and Asian Indians jointly (Figure 2B). Although a greater mean BMI was observed in CC subjects consuming a high saturated fat, the differences did not reach the statistical significance, and the interaction term between the *APOA2* genotype and saturated fat was not statistically significant ( $P=0.531$ ). Taking into account the low prevalence of the CC genotype in Chinese, and the possible heterogeneity of the effects compared with Asian Indians, we analyzed the *APOA2*-saturated fat interaction in Asian Indians. Although the interaction term was not statistically significant ( $P=0.445$ ), we observed similar interaction effects to those of the Mediterranean population. A higher saturated fat intake increased the differences in BMI between the *APOA2* genotypes (about 7.1% in the high saturated fat strata: 25.2 $\pm$ 0.5 Kg/m<sup>2</sup> in TT+TC versus 27.1 $\pm$ 1.2 Kg/m<sup>2</sup> in CC;  $P=0.155$  and less than 1% in the low saturated fat: 25.4 $\pm$ 0.6 Kg/m<sup>2</sup> in TT+TC versus 25.7 $\pm$ 1.5 Kg/m<sup>2</sup> in CC subjects;  $P=0.758$ ). These results could reflect a lack of

statistical power due to the small sample size (only 18 CC Indian subjects with dietary data). Considering our limitations in sample size in the Asian population for detecting as statistically significant the *APOA2*-saturated fat interaction in determining the continuous variable BMI, we focused on the dichotomic variable of obesity (BMI $\geq$ 27 Kg/m<sup>2</sup>). For this analysis, our sample size had sufficient power to detect statistically significant differences. Consistent with our previous results, we found that the CC genotype was only associated with a higher OR of obesity in subjects in the high-saturated fat stratum (Table 3). If saturated fat consumption was low, the CC genotype was not associated with obesity. These results were statistically significant for Asian Indians and also reached the statistical significance when the Chinese and Asian Indians were analyzed jointly (OR for obesity in CC homozygotes: 3.31; 95% CI: 1.08–10.18; P=0.036 in the high fat strata).

### Association between the *APOA2* –265T>C SNP, IR and diabetes. Modulation by saturated fat intake

In Asian Indians, the *APOA2* SNP was significantly associated with higher IR in the unadjusted analysis (3.7 $\pm$ 1.9 in CC vs 2.8 $\pm$ 2.8 in T-allele carriers; P=0.012). No statistically significant associations were found in Chinese (1.9 $\pm$ 1.3 in CC vs 1.8 $\pm$ 1.3 in T-allele carriers; P=0.611) or in Malays (2.0 $\pm$ 1.3 in CC vs 2.4 $\pm$ 2.3 in T-allele carriers; P=0.484). Next, we tested the *APOA2*-saturated fat interaction in determining IR in Chinese and Asian Indians (Figure 3) and found statistically significant results (P for interaction: 0.026). The CC genotype was associated with higher IR only when saturated fat intake was high ( $\geq$ 22g/d). The *APOA2*-saturated fat interaction remained statistically significant even after additional adjustment for BMI (P=0.039).

We further analyzed the association between the *APOA2* SNP and diabetes in Chinese and Asian Indians. In comparison with T-allele carriers, CC subjects had a higher risk of diabetes (OR: 2.14; 95% CI: 1.01–4.56, P=0.048) after adjustment for sex, age and ethnic group. However additional adjustment for smoking, drinking and BMI, attenuated the association (P=0.167). We tested if this association was modulated by saturated fat and found that, in the low saturated fat strata (<22 g/d), the CC genotype was not associated with higher diabetes risk (OR: 0.72; 95% CI: 0.08–6.72; P=0.770). However, in the high-saturated fat strata, the CC genotype was borderline (P=0.080) associated with higher diabetes risk (OR: 3.10; 95% CI: 0.87–11.02), after adjustment for sex, age, ethnic group, smoking, drinking and total energy intake. Further adjustment for other macronutrients and physical activity did not change the statistical significance of results.

In the Mediterranean population, we did not measure IR. In terms of diabetes, the *APOA2* SNP was not associated with diabetes in the whole population (P=0.927). However, in females, prevalence of diabetes was higher in CC women consuming less than 22g/d of saturated fat (P=0.045), suggesting a potential situation of reverse causation.

## DISCUSSION

We have previously reported in three American populations that the *APOA2* –265T>C SNP was associated with BMI or obesity only in the presence of high-saturated fat intake (11). Our current data are consistent with this gene-diet interaction and extends the findings to other geographical areas (Europe and Asia). When saturated fat intake is low (<22g/d), this SNP does not have any effect on BMI or obesity. However, when saturated fat intake is high ( $\geq$ 22g/d), significant differences in anthropometric variables were detected between CC individuals and T-allele carriers. Further adjustment for other macronutrients did not change the significance of our findings, supporting the specificity of saturated fat as a driver of this interaction.

Current recommendations to increase the consistency of gene-disease associations underline the importance of replication in independent samples (33). Our results, supporting the previous findings on anthropometric variables in the Framingham, the GOLDN and the Boston-Puerto Rican studies (11), contribute to increase the external validity of this gene-diet interaction. Moreover, we have reported for the first time a gene-diet interaction between the *APOA2* SNP and saturated fat intake in determining IR in Chinese and Asian Indians.

The mechanisms behind these novel interactions are still unknown; however, there are some lines of evidence supporting our findings. *APOA2*, the second most abundant apolipoprotein in HDL, plays a complex and relatively undefined role in lipoprotein metabolism, IR, obesity and atherosclerosis susceptibility (34). Castellani et al (12), found that *APOA2* transgenic mice had increased body-weight and body fat content in comparison to control mice. They also found that overexpression of *APOA2* in these mice increased IR as demonstrated by 2–3 fold higher fasting insulin levels and a delayed clearance of glucose bolus (12). These results were consistent with the previous observation of Breslow et al (35) in homozygous *APOA2* knockout mice. *APOA2* deficiency was associated with lower HDL, free fatty acid, glucose, and insulin levels, suggesting an insulin hypersensitivity state. More targeted mechanistic studies using these experimental models have confirmed the role of *APOA2* in IR (13,15,36). However, some controversial results regarding *APOA2* overexpression and IR and atherosclerosis have also been reported (34,37,38,39). One of the sources of discrepancies is that overexpression of human *APOA2* in mice does not always exert effects similar to those of murine *APOA2* (38). Interestingly, the other source of discrepancy and particularly relevant to our findings, is the demonstrated difference in *APOA2* expression depending on the type of diet administered (34,38). Thus, Escola-Gil et al (39) reported an *APOA2*-high fat diet interaction in determining the atherosclerotic effects of the *APOA2* transgenic mice. After an atherogenic high-fat (cocoa butter) diet, human *APOA2* transgenic mice developed more extensive aortic atherosclerosis than control mice. In contrast to the atherogenic diet-fed animals, there was no significant difference in the area of atherosclerotic lesion when they were fed a regular chow diet.

Taken together, the strong influence of a high-fat diet on the effects of the *APOA2* in transgenic animal models could help to explain our epidemiological gene-diet interaction. Although the –265T>C SNP has been associated with ~30% drop in basal transcription activity (19,20) and lower plasma *APOA2* concentrations (11) in standard conditions, a high-saturated fat diet could mask or reverse these effects resulting in a metabolic situation more favorable to increased body-weight and even IR. One possible hypothesis that we can suggest for integrating observations on animal and human models would be as follows: The C allele, associated in general with a lower *APOA2* expression would give rise to lower plasma concentrations of *APOA2*, found in our previous work (11). If the *APOA2* acted as a satiety signal, as is suggested for the *APOA4* (40), a lower plasma *APOA2* concentration would give rise to greater appetite. This appetite would preferably be for foods rich in saturated fat, and this higher fat intake would lead to greater weight and/or IR. In CC individuals who did not consume more saturated fat (maintaining it below 22 g/d), no weight gain or greater susceptibility to IR would be observed.

Previous human studies did not investigate the influence of diet on the effects of the –265T>C SNP, but controversial results on the association between this SNP and anthropometric measures have been reported. This SNP has been associated with lower waist circumference in men (19); with higher abdominal fat depots in women (22); with higher waist-to-hip ratio in French (23) and with higher BMI in White Americans (21). No association was also reported (41).

Type 2 diabetes has been linked to chromosome 1q21–24 in several populations (16–18), supporting a role for the APOA2 gene (located in 1q23) as a potential candidate; however, this association is still controversial. A recent case-control cohort comprising 3,093 French Caucasian subjects, did not find significant association between APOA2 SNPs and type 2 diabetes (23). However, significant gene-gene interactions with SNPs in the adjacent caldesmon 1 gene, so increasing diabetes risk, were found in Utah subjects (42). Likewise, Elbein et al (43) evaluated the association between APOA2 SNPs and diabetes in Northern European subjects. Although they did not find strong associations with diabetes, some variants (including the promoter polymorphism) were significantly associated with 2-h glucose and insulin secretion. However, none of these studies examined the potential modulation by dietary fat. We observed a statistically significant interaction of the –265T>C SNP with saturated fat intake in Chinese and Asian Indians. CC subjects had higher IR than T-allele carriers only with a high-saturated fat consumption. Similarly, we observed a higher OR for diabetes in the high-saturated fat strata (borderline significant). In the Mediterranean population, the APOA2 SNP was not associated with diabetes risk in the whole population. In the analysis stratified by saturated fat intake, we observed a higher prevalence of female diabetic subjects in the low saturated fat strata. Taking into account the characteristics of our high-risk Mediterranean population, in which the majority of individuals (>99%) with diabetes were previously diagnosed by their physician before the inclusion in the study, this observation could reflect a reverse causation situation due to dietary recommendations for treating diabetes, this being a limitation of our study. In contrast, the Asian population was a representative general population aged 18–69 years, in which a high percentage of individuals with diabetes (>55% of total) were newly detected. However, in the Asian population, the main limitation was the low prevalence of CC subjects in comparison with Caucasian populations. This observation, apart from affecting the statistical significance of the results, contributes to the fact that the population impact of this interaction in determining obesity and IR in Asians is low.

In conclusion, despite some sample size limitations and population specific results, we have replicated a gene-diet interaction involving the APOA2 –265T>C SNP and saturated fat intake in determining body-weight in a Mediterranean and an Asian population, so increasing the external validity of our previously reported results in American populations (11). In addition, our results suggest that this interaction may also play a role in IR and diabetes. However, prospective studies are required to better characterize this dietary modulation.

## Acknowledgments

**Funding/Support:** This work was supported by National Heart, Lung, and Blood Institute grants U 01 HL72524 and HL-54776, National Institute of Diabetes and Digestive and Kidney Diseases, Grant Number DK075030 and by contracts 53-K06-5-10 and 58-1950-9-001 from the US Department of Agriculture Research Service and grant from the Ministerio de Ciencia e Innovación (PR2008-268 and PR2009-392), the Generalitat Valenciana (GVACOMP2010-181), the Centro Nacional de Investigaciones Cardiovasculares (CNIC06) and the Instituto de Salud Carlos III (CIBER CB06/03/0035, RD07/0067/0006 and PI070954), Spain.

## REFERENCES


1. Rankinen T, Zuberi A, Chagnon YC, Weisnagel SJ, Argyropoulos G, Walts B, et al. The human obesity gene map: the 2005 update. *Obesity (Silver Spring)*. 2006; 14:529–644. [PubMed: 16741264]
2. Walley AJ, Asher JE, Froguel P. The genetic contribution to non-syndromic human obesity. *Nat Rev Genet*. 2009; 10:431–442. [PubMed: 19506576]



3. Thorleifsson G, Walters GB, Gudbjartsson DF, Steinthorsdottir V, Sulem P, Helgadóttir A, et al. Genome-wide association yields new sequence variants at seven loci that associate with measures of obesity. *Nat Genet.* 2009; 41:18–24. [PubMed: 19079260]
4. Willer CJ, Speliotes EK, Loos RJ, Li S, Lindgren CM, Heid IM, et al. Six new loci associated with body mass index highlight a neuronal influence on body weight regulation. *Nat Genet.* 2009; 41:25–34. [PubMed: 19079261]
5. Frayling TM, Timpson NJ, Weedon MN, Zeggini E, Freathy RM, Lindgren CM, et al. A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. *Science.* 2007; 316:889–894. [PubMed: 17434869]
6. Dina C, Meyre D, Gallina S, Durand E, Körner A, Jacobson P, et al. Variation in FTO contributes to childhood obesity and severe adult obesity. *Nat Genet.* 2007; 39:724–726. [PubMed: 17496892]
7. Bouchard C. Gene-environment interactions in the etiology of obesity: defining the fundamentals. *Obesity (Silver Spring).* 2008; 16:S5–S10. [PubMed: 19037213]
8. Andreasen CH, Stender-Petersen KL, Mogensen MS, Torkov SS, Wegner L. Low physical activity accentuates the effect of the FTO rs9939609 polymorphism on body fat accumulation. *Diabetes.* 2008; 57:95–101. [PubMed: 17942823]
9. Rampersaud E, Mitchell BD, Pollin TI, Fu M, Shen H, O'Connell JR, et al. Physical activity and the association of common FTO gene variants with body mass index and obesity. *Arch Intern Med.* 2008; 168:1791–1797. [PubMed: 18779467]
10. Vimalaswaran KS, Li S, Zhao JH, Luan J, Bingham SA, Khaw KT, et al. Physical activity attenuates the body mass index-increasing influence of genetic variation in the FTO gene. *Am J Clin Nutr.* 2009; 90:425–428. [PubMed: 19553294]
11. Corella D, Peloso G, Arnett DK, Demissie S, Cupples LA, Tucker K, et al. APOA2, dietary fat, and body mass index replication of a gene-diet interaction in 3 independent populations. *Arch Internal Med.* 2009; 169:1897–1906. [PubMed: 19901143]
12. Castellani LW, Goto AM, Lusic AJ. Studies with apolipoprotein A-II transgenic mice indicate a role for HDLs in adiposity and insulin resistance. *Diabetes.* 2001; 50:643–651. [PubMed: 11246886]
13. Castellani LW, Gargalovic P, Febbraio M, Charugundla S, Jien ML, Lusic AJ. Mechanisms mediating insulin resistance in transgenic mice overexpressing mouse apolipoprotein A-II. *J Lipid Res.* 2004; 45:2377–2387. [PubMed: 15466364]
14. Suto J. Quantitative trait locus analysis of plasma cholesterol levels and body weight by controlling the effects of the ApoA2 allele in mice. *J Vet Med Sci.* 2007; 69:385–392. [PubMed: 17485926]
15. Castellani LW, Nguyen CN, Charugundla S, Weinstein MM, Doan CX, Blaner WS, et al. Apolipoprotein AII is a regulator of very low density lipoprotein metabolism and insulin resistance. *J Biol Chem.* 2008; 283:11633–11644. [PubMed: 18160395]
16. Elbein SC, Hoffman MD, Teng K, Leppert MF, Hasstedt SJ. A genome-wide search for type 2 diabetes susceptibility genes in Utah Caucasians. *Diabetes.* 1999; 48:1175–1182. [PubMed: 10331426]
17. Vionnet N, Hani EH, Dupont S, Gallina S, Francke S, Dotte S, et al. Genomewide search for type 2 diabetes-susceptibility genes in French whites: evidence for a novel susceptibility locus for early-onset diabetes on chromosome 3q27-qter and independent replication of a type 2-diabetes locus on chromosome 1q21–q24. *Am J Hum Genet.* 2000; 67:1470–1480. [PubMed: 11067779]
18. Xiang K, Wang Y, Zheng T, Jia W, Li J, Chen L, et al. Genome-wide search for type 2 diabetes/ impaired glucose homeostasis susceptibility genes in the Chinese: significant linkage to chromosome 6q21–q23 and chromosome 1q21–q24. *Diabetes.* 2004; 53:228–234. [PubMed: 14693720]
19. van 't Hooft FM, Ruotolo G, Boquist S, de Faire U, Eggertsen G, Hamsten A. Human evidence that the apolipoprotein A-II gene is implicated in visceral fat accumulation and metabolism of triglyceride-rich lipoproteins. *Circulation.* 2001; 104:1223–1228. [PubMed: 11551871]
20. Takada D, Emi M, Ezura Y, Nobe Y, Kawamura K, Iino Y. Interaction between the LDL-receptor gene bearing a novel mutation and a variant in the apolipoprotein A-II promoter: molecular study in a 1135-member familial hypercholesterolemia kindred. *J Hum Genet.* 2002; 47:656–664. [PubMed: 12522687]


21. Corella D, Arnett DK, Tsai MY, Kabagambe EK, Peacock JM, Hixson JE, et al. The -256T>C polymorphism in the apolipoprotein A-II gene promoter is associated with body mass index and food intake in the genetics of lipid lowering drugs and diet network study. *Clin Chem*. 2007; 53:1144–1152. [PubMed: 17446329]
22. Lara-Castro C, Hunter GR, Lovejoy JC, Gower BA, Fernandez JR. Apolipoprotein A-II polymorphism and visceral adiposity in African-American and white women. *Obesity*. 2005; 13:507–512.
23. Duesing K, Charpentier G, Marre M, Tichet J, Hercberg S, Balkau B, et al. Evaluating the association of common APOA2 variants with type 2 diabetes. *BMC Med Genet*. 2009; 10:13. [PubMed: 19216768]
24. Estruch R, Martínez-González MA, Corella D, Salas-Salvadó J, Ruiz-Gutiérrez V, Covas MI, et al. Effects of a Mediterranean-style diet on cardiovascular risk factors: a randomized trial. *Ann Intern Med*. 2006; 145:1–11. [PubMed: 16818923]
25. Cutter J, Tan BY, Chew SK. Levels of cardiovascular disease risk factors in Singapore following a national intervention programme. *Bull World Health Organ*. 2001; 79:908–915. [PubMed: 11693972]
26. Deurenberg-Yap M, Chew SK, Lin VF, Tan BY, van Staveren WA, Deurenberg P. Relationships between indices of obesity and its co-morbidities in multi-ethnic Singapore. *Int J Obes Relat Metab Disord*. 2001; 25:1554–1562. [PubMed: 11673781]
27. Deurenberg P, Deurenberg-Yap M, Guricci S. Asians are different from Caucasians and from each other in their body mass index/body fat per cent relationship. *Obes Rev*. 2002; 3:141–146. [PubMed: 12164465]
28. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*. 1985; 28:412–419. [PubMed: 3899825]
29. Alberti, KG.; Zimmet, PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1. Diagnosis and classification of diabetes mellitus provisional report of a WHO consultation; *Diabet Med*. 1998. p. 539-553.
30. Elosua R, Garcia M, Aguilar A, Molina L, Covas MI, Marrugat J. Investigators of the MARATDON Group. Validation of the Minnesota Leisure Time Physical Activity Questionnaire In Spanish Women. *Med Sci Sports Exerc*. 2000; 32:1431–1437. [PubMed: 10949009]
31. Martin-Moreno JM, Boyle P, Gorgojo L, Maisonneuve P, Fernandez-Rodriguez JC, Salvini S, et al. Development and validation of a food frequency questionnaire in Spain. *Int J Epidemiol*. 1993; 22:512–519. [PubMed: 8359969]
32. Deurenberg-Yap M, Li T, Tan WL, Tan WL, van Staveren WA, Chew SK, et al. Can dietary factors explain differences in serum cholesterol profiles among different ethnic groups (Chinese, Malays and Indians) in Singapore? *Asia Pac J Clin Nutr*. 2001; 10:39–45. [PubMed: 11708607]
33. Little J, Higgins JP, Ioannidis JP, Moher D, Gagnon F, von Elm E, et al. STrengthening the REporting of Genetic Association studies (STREGA): an extension of the STROBE Statement. *Ann Intern Med*. 2009; 150:206–215. [PubMed: 19189911]
34. Martín-Campos JM, Escolà-Gil JC, Ribas V, Blanco-Vaca F. Apolipoprotein A-II, genetic variation on chromosome 1q21–q24, and disease susceptibility. *Curr Opin Lipidol*. 2004; 15:247–253. [PubMed: 15166779]
35. Weng W, Breslow JL. Dramatically decreased high density lipoprotein cholesterol, increased remnant clearance, and insulin hypersensitivity in apolipoprotein A-II knockout mice suggest a complex role for apolipoprotein A-II in atherosclerosis susceptibility. *Proc Nat Acad Sci USA*. 1996; 93:14788–14794. [PubMed: 8962133]
36. Sauvaget D, Chauffeton V, Dugué-Pujol S, Kalopissis AD, Guillet-Deniau I, Foufelle F, et al. In vitro transcriptional induction of the human apolipoprotein A-II gene by glucose. *Diabetes*. 2004; 53:672–678. [PubMed: 14988251]
37. Escolà-Gil JC, Blanco-Vaca F, Julve J. Overexpression of human apolipoprotein A-II in transgenic mice does not increase their susceptibility to insulin resistance and obesity. *Diabetologia*. 2002; 45:600–601. [PubMed: 12032642]

38. Blanco-Vaca F, Escolà-Gil JC, Martín-Campos JM, Julve J. Role of apoA-II in lipid metabolism and atherosclerosis: advances in the study of an enigmatic protein. *J Lipid Res.* 2001; 42:1727–1739. [PubMed: 11714842]
39. Escola-Gil JC, Marzal-Casacuberta A, Julve-Gil J, Ishida BY, Ordonez-Llanos J, Chan L, et al. Human apolipoprotein A-II is a pro-atherogenic molecule when it is expressed in transgenic mice at a level similar to that in humans: evidence of a potentially relevant species-specific interaction with diet. *J Lipid Res.* 1998; 39:457–462. [PubMed: 9580110]
40. Culnan DM, Cooney RN, Stanley B, Lynch CJ. Apolipoprotein A-IV, a putative satiety/antiatherogenic factor, rises after gastric bypass. *Obesity (Silver Spring).* 2009; 17:46–52. [PubMed: 18948973]
41. Xiao J, Zhang F, Wiltshire S, Hung J, Jennens M, Beilby JP, et al. The apolipoprotein AII rs5082 variant is associated with reduced risk of coronary artery disease in an Australian male population. *Atherosclerosis.* 2008; 199:333–339. [PubMed: 18179799]
42. Hasstedt SJ, Chu WS, Das SK, Wang H, Elbein SC. Type 2 diabetes susceptibility genes on chromosome 1q21–24. *Ann Hum Genet.* 2008; 72:163–169. [PubMed: 18269685]
43. Elbein SC, Chu W, Ren Q, Wang H, Hemphill C, Hasstedt SJ. Evaluation of apolipoprotein A-II as a positional candidate gene for familial Type II diabetes, altered lipid concentrations, and insulin resistance. *Diabetologia.* 2002; 45:1026–1033. [PubMed: 12136402]




**Figure 1.**

Association between the *APOA2* -265T>C polymorphism and body-weight in **(A)** Mediterranean subjects (n= 374 TT, 398 TC and 135 CC) and in **(B)** Asian Indians (n= 305 TT, 160 TC and 29 CC). Adjusted means and error bars by genotype. Means were adjusted for sex, age, tobacco smoking and alcohol consumption. \* P values for mean comparison between CC and T allele carriers obtained in the multivariate adjusted models. Error bars: SE of means.



**Figure 2.**

Interaction between the *APOA2*  $-265T>C$  polymorphism and saturated fat intake on BMI in (A) Mediterranean subjects ( $n=772$  T-allele carriers and  $n=135$  CC) and in (B) Chinese and Asian Indians ( $n=1483$  T-allele carriers and 28 CC subjects). Adjusted means of BMI are shown depending on the *APOA2*  $-265T>C$  polymorphism according to the strata of saturated fat intake (below and above 22g/d). Means were adjusted for sex, age (as continuous), tobacco smoking (as categorical), alcohol consumption (as categorical), diabetes status (as categorical), lipid medication (only in the Mediterranean population), ethnicity (in the Asian population) and total energy intake (as continuous). P values for the interaction terms between saturated fat intake (as dichotomous) and the *APOA2* polymorphism in each population were obtained in the hierarchical multivariate interaction model. In the stratified analysis by saturated fat intake levels, P values for mean comparisons of BMI between *APOA2* genotypes were estimated after multivariate adjustment for the covariates indicated above. Bars indicate standard error (SE) of means.



**Figure 3.**

Interaction between the APOA2 -265T>C polymorphism and saturated fat intake in determining IR in Chinese (n=1204) and Asian Indians (n=307). Adjusted means of IR are shown depending on the APOA2 -265T>C polymorphism (n=1493 T-allele carriers and 28 CC) according to the strata of saturated fat intake (below and above 22g/d). Means were adjusted for sex, age (as continuous), ethnicity, tobacco smoking (as categorical), alcohol consumption (as categorical), and total energy intake (as continuous). P values for the interaction terms between saturated fat intake (as dichotomous) and the APOA2 polymorphism in each population were obtained in the hierarchical multivariate interaction model. In the stratified analysis by saturated fat intake levels, P values for mean comparisons of BMI between APOA2 genotypes were estimated after multivariate adjustment for the covariates indicated above. All statistical analyses were carried out with the ln transformed IR. Bars indicate standard error (SE) of means.

**Table 1**

General characteristics of the studied populations

	Mediterranean	Multiethnic Asian population		
	population (n=907)	Chinese (n=2506)	Malays (n=695)	Asian Indians (n=494)
	Mean SD	Mean SD	Mean SD	Mean SD
Male/Female	325/582	1143/1363	335/360	236/258
Age (years)	67 (6)	38 (12)	39 (13)	40 (12)
Weight (kg)	77.5 (13.1)	60.5 (12.3)	65.8 (13.5)	66.4 (13.0)
BMI (Kg/m <sup>2</sup> )	30.9 (5.1)	22.7 (3.7)	25.6 (5.0)	25.1 (4.6)
Waist (cm)	104 (12)	78 (11)	83 (12)	85 (12)
HDL-C (mg/dL)	53.2 (14.1)	55.4 (14.8)	50.3 (12.9)	44.1 (11.3)
Fasting triglycerides (mg/dL)	130.9 (80.9)	124.6 (106.8)	148.6 (101.5)	151.3 (127.3)
Fasting glucose (mg/dL)	122.7 (40.2)	100.8 (23.4)	109.8 (37.8)	113.4 (39.6)
HOMA-IR	ND	1.8 (1.3)	2.4 (2.2)	2.9 (2.8)
Energy intake (Kcal/d)*	2253 (669)	2045 (744)	2177 (931)	2220 (771)
Total fat intake (g/d)*	97.4 (32.9)	61.8 (29.3)	69.2 (39.5)	69.2 (29.3)
Total fat intake (% energy)*	39.1 (7.1)	26.6 (5.3)	27.9 (6.0)	27.7 (5.6)
Saturated fat intake (g/d)*	25.7 (10.0)	24.2 (12.7)	29.8 (17.8)	28.9 (15.2)
Saturated fat (% energy)*	10.2 (2.3)	10.3 (2.5)	11.8 (3.2)	11.4 (2.9)
Monounsaturated fat (% energy)*	19.2 (4.8)	9.2 (2.3)	9.0 (2.5)	8.1 (2.2)
Polyunsaturated fat (% energy)*	6.3 (2.3)	5.2 (2.0)	4.8 (2.0)	5.8 (2.6)
Saturated fat < 22g/d*; n (%)	362 (39.9)	614 (51.0)	165 (40.0)	115 (37.5)
Drinkers; n (%)	467 (51.5)	1132 (45.2)	55 (7.9)	162 (32.8)
Current smokers; n (%)	106 (11.7)	316 (12.6)	162 (23.3)	73 (14.8)
BMI>=30 Kg/m <sup>2</sup>	478 (52.7)	93 (3.7)	112 (16.1)	61 (12.3)
BMI>=27 Kg/m <sup>2</sup>	711 (78.4)	310 (12.4)	234 (33.7)	153 (31.0)
Diabetes; n (%)	443 (48.8)	182 (7.3)	103 (14.8)	97 (19.6)
Hypertension; n (%)	724 (79.8)	209 (8.3)	62 (8.9)	56 (11.3)
APOA2-265T>C genotype; n (%)				
TT	374 (41.2)	2080 (83.0)	558 (80.3)	305 (61.7)
TC	398 (43.9)	405 (16.2)	128 (18.4)	160 (32.4)
CC	135 (14.9)	21 (0.8)	9 (1.3)	29 (5.9)

\* Dietary variables were only determined in a random sample (1 in 2) of the Asian population (n=1,923 subjects: 1204 Chinese, 412 Malays and 307 Asian Indians)

HOMA-IR: Homeostasis model assessment of insulin resistance

ND: Not determined

**Table 2**  
Association of the APOA2 polymorphism with anthropometric variables in the Mediterranean and the Asian populations

	Mediterranean population			Asian Indians			Chinese			Malays		
	TT+TC (n=772)	CC (n=135)	p <sup>1</sup>	TT+TC (n=465)	CC (n=29)	p <sup>2</sup>	TT+TC (n=2485)	CC (n=21)	p <sup>2</sup>	TT+TC (n=686)	CC (n=9)	p <sup>2</sup>
	Mean	SD		Mean	SD		Mean	SD		Mean	SD	
Weight (kg)	77.0	(12.8)	0.047	65.9	(12.2)	0.018	60.5	(12.3)	0.021	65.8	(13.6)	0.951
BMI (Kg/m <sup>2</sup> )	30.6	(5.1)	0.049	24.9	(4.6)	0.038	22.7	(3.7)	0.006	25.6	(5.0)	0.759
Waist (cm)	103.8	(12.1)	0.605	84.9	(11.8)	0.375	78.1	(10.7)	0.067	82.8	(12.1)	0.987
Obesity <sup>*</sup> ; n (%)	404	(52.3)	0.594	136	(29.2)	0.264	308	(12.4)	0.004	231	(33.7)	0.590

<sup>1</sup> P values for the comparison of unadjusted means (for continuous variables) or percentages (for categorical variables) between APOA2 genotypes

<sup>2</sup> P values obtained in the multivariate adjusted (for sex, age, tobacco smoking, alcohol and diabetes) models

\* Obesity was defined as BMI $\geq$  30 Kg/m<sup>2</sup> in the Mediterranean and as BMI $\geq$ 27 Kg/m<sup>2</sup> in the Asian population

SD: Standard deviation



Table 3

Associations between the APOA2 -265C>T polymorphism and obesity depending on the saturated fat intake category (lower or higher than 22 g/d) in the Asian population. Multivariate logistic regression analysis

	Saturated fat	APOA2 Genotype	Obesity* (n)		OR <sup>1</sup>	(95% CI)	P <sup>1</sup>	OR <sup>2</sup>	(95% CI)	P <sup>2</sup>
			No	Yes						
Asian Indians	<22g/d	TT+TC	78	30	1			1		
		CC	5	2	0.913	(0.151–5.525)	0.921	1.048	(0.158–6.956)	0.962
	>=22g/d	TT+TC	132	49	1			1		
		CC	4	7	4.502	(1.085–18.579)	0.038	4.834	(1.172–19.940)	0.029
Chinese	<22g/d	TT+TC	540	68	1			1		
		CC	6	0	0.000	-	0.493	0.000	-	0.999
	>=22g/d	TT+TC	505	81	1			1		
		CC	3	1	2.141	(0.207–22.180)	0.523	2.009	(0.186–21.677)	0.565
Chinese and Asian Indians**	<22g/d	TT+TC	618	98	1			1		
		CC	11	2	0.557	(0.101–3.063)	0.501	0.599	(0.101–3.546)	0.572
	>=22g/d	TT+TC	637	130	1			1		
		CC	7	8	3.310	(1.076–10.182)	0.036	3.332	(1.074–10.337)	0.037

\* BMI>=27 Kg/m<sup>2</sup>

<sup>1</sup> Models were adjusted for sex, age, tobacco smoking, alcohol consumption and diabetes

<sup>2</sup> Model were adjusted for sex, age, tobacco smoking alcohol consumption, diabetes, total energy intake, total carbohydrates and proteins

\*\* Additional adjustment for ethnic group was carried out