



---

**Título artículo / Títol article:** Sex pheromones are not always attractive: changes induced by learning and illness in mice

**Autores / Autors** E. Lanuza, A. Martín Sánchez, P. Marco Monclús, B. Cádiz-Moretti, L. Fortes Marco, A. Hernández Hernández, L. McLean, R. J. Beynon, J. L. Hurst, F. Martínez-García

**Revista:** Animal Behavior, v. 97 (2014)

**Versión / Versió:** Preprint de l'autor

**Cita bibliográfica / Cita bibliogràfica (ISO 690):** LANUZA, E...[et al.] Sex pheromones are not always attractive: changes induced by learning and illness in mice. Animal Behavior, v. 97 (2014), pp. 265-272

**url Repositori UJI:** <http://repositori.uji.es/xmlui/handle/10234/124352>

---

1 **TITLE: Sex pheromones are not always attractive: changes induced by learning,**  
2 **puberty, lactation and illness in mice**

3 **AUTHORS:** Enrique Lanuza<sup>1\*</sup>, Ana Martín-Sánchez<sup>1</sup>, Pau Marco-Manclús<sup>1</sup>, Bernardita  
4 Cádiz-Moretti<sup>1</sup>, Lluís Fortes-Marco<sup>1</sup>, Adoración Hernández-Martínez<sup>1</sup>, Lynn McLean<sup>2</sup>,  
5 Robert J. Beynon<sup>2</sup>, Jane L. Hurst<sup>2</sup>, and Fernando Martínez-García<sup>1</sup>

6 1) Laboratory of Functional and Comparative Neuroanatomy, Depts. of Biologia  
7 Cel·lular and Biologia Funcional, Facultat de Ciències Biològiques, Universitat  
8 de València, Spain.

9 2) Institute of Integrative Biology, University of Liverpool, UK.

10 \*Correspondence to: Enrique Lanuza. Dept. de Biologia Cel·lular. Facultat de Ciències  
11 Biològiques. Universitat de València. Carrer Dr. Moliner, 50. ES-46100, Burjassot. València.  
12 Spain.

13 Phone: 34 96 354 33 83 Fax: 34 96 354 34 04.

14 E-mail: Enrique.Lanuza@uv.es

15 NUMBER OF WORDS (excluding references and abstract): 5803

16 NUMBER OF FIGURES: 2

17

18 ABSTRACT

19 A male-specific major urinary protein named darcin is attractive to female mice, stimulates a  
20 learned attraction to volatile components of a male's urinary odour (Roberts et al., 2010,  
21 BMC Biol 8:75) and induces spatial learning (Roberts et al., 2012, Science 338:1462-5). In  
22 the present work we show that darcin also induces a learned attraction for a previously  
23 neutral olfactory stimulus (the odorant isoamyl acetate). However, the attractive properties of  
24 darcin may change as a function of female physiological state. For example, during the  
25 period of lactation female mice display aggressive behaviour against intruders, which is  
26 enhanced when confronted with adult males. Therefore, the endocrinological status of the  
27 females radically changes the behavioural response. The situation at puberty is somewhat  
28 similar: prepubertal females avoid adult male chemical signals, whereas post-pubertal  
29 females show attraction for these same signals. In addition, we report another situation in  
30 which the presence of darcin is not attractive to adult female mice. Urine of males parasitized  
31 by the intestinal nematode *Aspiculuris tetraptera* shows no attractive value for female mice,  
32 despite apparently normal presence of darcin. In this case, the loss of the attractive value is  
33 not due to physiological changes in the receptor females but is likely to be due to the  
34 presence of unknown signals of infection whose detection overrides the attraction normally  
35 induced by darcin.

36

37 KEY WORDS: vomeronasal, olfactory, sexual attraction, learning, maternal aggression,  
38 puberty, illness cues.

39

40 Pheromones were originally defined as “substances which are secreted to the outside by  
41 an individual and received by a second individual of the same species, in which they  
42 release a specific reaction, for example, a definite behavior or a developmental process”  
43 (Karlson & Luscher, 1959). Although this definition has been very useful for more than  
44 50 years (Wyatt, 2009), the response to pheromones (at least in mammals) may vary  
45 depending on a number of factors, including the previous experience or the hormonal  
46 status of the receiver (Wyatt, 2010). Moreover, the behavioural response elicited by  
47 pheromones may also be observed as learned responses to stimuli previously associated  
48 with the pheromones, so that these stimuli acquire pheromone-like properties (Moncho-  
49 Bogani, Lanuza, Hernández, Novejarque, & Martínez-García, 2002). Here, we review  
50 relevant examples in mice, in which the attractive value of male sexual pheromones is  
51 transferred to a neutral stimulus or changes either as a function of the hormonal  
52 condition of the receiver females or the health status of scent donor males. In addition,  
53 we include data from two experiments giving support to the ideas presented in this  
54 review.

55

### 56 **Pheromone-induced olfactory learning**

57 Odours are easily associated with either positive or negative experiences, becoming  
58 secondary attractive or aversive stimuli that strongly influence behavioural responses in  
59 many mammalian species, including humans (Herz & Cupchik, 2005). In rodents,  
60 olfactory stimuli play a key role in many aspects of socio-sexual behaviours (Brennan &  
61 Kendrick, 2006). The experience with social chemical signals influences later  
62 responses. For example, chemical signals present in urine of male mice, detected  
63 through the vomeronasal system, have reinforcing properties able to induce appetitive  
64 associative learning (Martínez-Ricós, Agustín-Pavón, Lanuza, & Martínez-García,

65 2007, 2008; Moncho-Bogani et al., 2002; Ramm, Cheetham, & Hurst, 2008), in such a  
66 way that other volatiles present in urine may become secondary attractive odorants  
67 (Martínez-García et al., 2009). More recently, a male-specific urinary protein named  
68 darcin, has been shown to be able to induce this kind of olfactory learning (Roberts et  
69 al., 2010) and also spatial learning (Roberts, Davidson, McLean, Beynon, & Hurst,  
70 2012). We hypothesized that darcin would be able to induce a secondary attraction by  
71 association with a neutral odorant (not present in urine). To test this possibility, we  
72 performed the following experiment.

73 EXPERIMENT 1: Inducing attraction for a neutral odorant by association with darcin

74 Material and Methods

75 *Subjects*

76 For the present study, 15 adult female mice (12-16 weeks) of the CD1 outbred strain  
77 were used (Janvier Labs., Le Genest-Saint-Isle, Saint-Berthevin Cedex, France).

78 Treatment of the animals employed in the experiments reported in this paper complied  
79 with the European Union Council Directive of June 3<sup>rd</sup>, 2010 (6106/1/10 REV1),  
80 according to which procedures were approved by the Committee of Ethics on Animal  
81 Experimentation of the University of Valencia (protocol number A1283764105250).

82 Procedures also adhered to the ASAB/ABS Guidelines for the Use of Animals in  
83 Research.

84 The females were sexually naïve and had never been exposed to chemical signals from  
85 sexually mature males. To achieve this, pregnant females were housed in a clean room  
86 without males, in standard macrolon transparent cages with a wire lid (21.5 x 46.5 x  
87 14.5 cm, ref. 1000, Panlab, Barcelona, Spain) filled with soft wood bedding (Souralit  
88 S.L., ref. 3000, Barcelona, Spain), provided with nesting material (shredded paper) and  
89 enriched with cardboard tubes. The room was maintained at 22-24 °C, 60–80% RH and

90 a 12:12 h light:dark cycle, with lights on at 0800 hours). Food (Teklad Global 14%  
91 Protein Rodent Maintenance Diet, Harlan, ref: 2014) and water were available *ad*  
92 *libitum*. Nineteen days after delivery (early before puberty), pups were sexed and males  
93 were removed. Their female siblings were brought to a clean room in complete absence  
94 of adult male chemical signals, where they were kept in groups of 5-6 per cage (the  
95 stock housing conditions in the experimental room were the same as described above  
96 for the pregnant females). Food and water were available *ad libitum* except during the  
97 preference tests (five-minute long) and the training sessions (15-minute long). Welfare  
98 assessment took place during cage cleaning, and included non-invasive indicators. In  
99 the neonates, skin colour, activity and presence of the milk spot were observed; at  
100 weaning and in the adult, general appearance, size, coat condition, posture, gait, activity  
101 levels, interaction with the environment and clinical signs were observed (Wells et al.,  
102 2006). After weaning, animals were only manipulated for cage cleaning once a week.  
103 Since general appearance and size were evaluated as normal, no further care was  
104 necessary. Mice were handle following the standard practice of picking them up by  
105 gently holding the base of the tail and helping them onto the handler's arm, avoiding to  
106 hold them in the air. All procedures involved in this study were non-invasive  
107 behavioural tests. At the end of the experiments, animals were euthanized with an  
108 intraperitoneal overdose of sodium pentobarbital (92 mg/kg), as indicated in the  
109 approved protocol (cited above). The male siblings were either used for anatomical  
110 studies (protocols approved by the Committee of Ethics on Animal Experimentation of  
111 the University of Valencia, under the same reference number A1283764105250;  
112 published elsewhere, Otero-García et al., 2014) or euthanized as described above.  
113 *Stimuli*

114 We chose two odorants that have been used frequently as olfactory stimuli in the  
115 literature: isoamyl acetate (e.g., Angely & Coppola, 2010; Panreac, Barcelona, Spain)  
116 and citralva (e.g., Martínez-Ricós et al., 2007; geranonitrile, 3,7-dimethyl-2,6-  
117 octadiene-1-nitrile, kindly provided by International Flavours and Fragrances, Ventos,  
118 Barcelona, Spain). Both odorants were diluted 1:1000 in phosphate buffer (0.01 M) with  
119 0.01% Triton X-100. In a pilot test of olfactory preference we proved that isoamyl  
120 acetate and citralva were investigated equally.

### 121 *Preference tests*

122 Animals were habituated to handling and to the test cage over 3 days, 10 min per day,  
123 between 15:00 and 20:00 hours. Preference tests were performed in 25 x 50 x 30 cm  
124 cages. A 4 x 4 cm piece of filter paper impregnated with 5 µl of one of the stimulus  
125 odorants was presented in each opposing side of the cage. These impregnated papers  
126 were fixed to the bottom of the cage with a metallic cover, leaving exposed a circular  
127 area (diameter = 3.5 cm) that allowed direct nasal contact with the paper but prevented  
128 the animals to gnaw or remove it.

129 For this olfactory preference test (citralva vs isoamyl acetate) females were released in  
130 the centre of the cage, the experimenter left the room, and the behaviour was videotaped  
131 for 5 min. The time that the animal spent in the circular area covered by the paper was  
132 measured by tracking animals automatically using the video analyser software Smart  
133 2.5 (Panlab, Barcelona, Spain; see Fig. 1A). Since we observed that the animals lost  
134 interest in the olfactory stimuli at the end of the test, we restricted the analysis of the  
135 data to the first four minutes.

136 Following the pre-training preference test, the females were run in a second test in  
137 which the isoamyl acetate-scented paper was also impregnated with 8 µl of recombinant  
138 darcin (r-darcin, diluted 1.1 µg/µl, Roberts et al., 2010). Over the next four days, four

139 training sessions (one per day) were performed in which a piece of paper (not fixed)  
140 scented with 5  $\mu$ l of isoamyl acetate and 8  $\mu$ l of r-darcin was presented to the females  
141 during 15 min/day in the centre of a different cage (29 x 15 x 29.8 cm), and thus in a  
142 different context to that of the preference tests. Finally, a post-training olfactory  
143 preference test was performed (citalva vs isoamyl acetate) identical to the pre-training  
144 test described above. The location (left or right) of the citralva and isoamyl acetate-  
145 scented papers was decided randomly at the beginning of the experiment and kept fixed  
146 for all the animals and tests.

147 After finishing this experiment, we wanted to discard the possibility that repeated  
148 presentations of the odorant, by themselves, induced preference for this familiar  
149 stimulus (against the other odorant, which is less familiar). To test this idea, we  
150 performed a second (control) experiment (n = 12, 16 weeks of age, Janvier Labs., Saint-  
151 Berthevin Cedex, France) identical to the previous one, except for the absence of darcin  
152 in the preference test and training sessions. Since females of this control experiment  
153 were not going to be exposed to male-derived chemosignals, they were acquired from  
154 Janvier as young adults, without preventing their prepubertal exposure to male odours.  
155 Housing and care conditions were the same as described above.

#### 156 *Statistical analysis*

157 The time spent in the area occupied by the scented paper was analysed with a repeated-  
158 measures ANOVA with the TEST (control, 1<sup>st</sup> odour preference test; darcin preference  
159 test; 2<sup>nd</sup> odour preference test) and SIDE (citalva vs. isoamyl acetate) as intra-subjects  
160 factors. The normality of the data was previously confirmed with a Kolmogorov-  
161 Smirnov test with Lillieford's correction. Analyses were performed with the SPSS 15.0  
162 software package.

#### 163 Results



164 The results of the repeated measures ANOVA for the time spent within each of the  
165 circular areas (SIDE) during the four TESTS (Control; 1<sup>st</sup> odour preference test; darcin  
166 preference test; 2<sup>nd</sup> odour preference test; Figure 1B) revealed significant main effects  
167 of SIDE ( $F_{1,14} = 7.87$ ,  $p = 0.014$ ), and non-significant main effect of TEST or the SIDE  
168 x TEST interaction ( $F < 1$ ,  $p > 0.6$  in both cases). The analysis of the simple effects of  
169 the factor SIDE in each test showed that both areas were equally investigated in the  
170 control (clean versus clean,  $F < 1$ ,  $p > 0.4$ ) and the 1<sup>st</sup> odour preference test (citalva vs.  
171 isoamyl acetate,  $F_{1,14} = 1.18$ ,  $p = 0.16$ ). By contrast, the females spent significantly  
172 more time in the area occupied by the paper scented with isoamyl acetate plus darcin  
173 ( $F_{1,14} = 5.07$ ,  $p = 0.032$ ). Finally, in the 2<sup>nd</sup> (post-training) olfactory preference test  
174 (citalva vs. isoamyl acetate), females again spent more time in the area occupied by the  
175 isoamyl acetate-scented paper ( $F_{1,14} = 5.77$ ,  $p = 0.031$ ).

176 In the control group, in which the same procedure was used but no darcin was present,  
177 the results of the repeated measures ANOVA for the time spent within each of the  
178 circular areas (SIDE) during the four TESTS (Figure 1B) showed no significant main  
179 effects of SIDE ( $F_{1,11} < 1$ ,  $p = 0.51$ ), TEST ( $F_{3,9} < 1$ ,  $p = 0.71$ ), or their interaction ( $F_{3,9}$   
180  $= 2.06$ ,  $p = 0.17$ ). The analysis of the simple effects of the factor SIDE in each test  
181 showed that both areas were equally investigated in all cases (clean versus clean:  $F_{1,11} =$   
182  $3.23$ ,  $p = 0.1$ ; first odour preference test:  $F_{1,11} = 1.02$ ,  $p = 0.33$ ; second odour preference  
183 test:  $F_{1,11} < 1$ ,  $p = 0.99$ ; third odour preference test:  $F_{1,11} = 1.37$ ,  $p = 0.26$ ).

#### 184 Discussion

185 The results of the present experiments show that a neutral odorant, such as isoamyl  
186 acetate, which is not significantly preferred by female mice, becomes a preferred  
187 olfactory stimulus by presenting it together with the sexual pheromone darcin. The  
188 repeated presentation of the odorant, as shown by the control experiment, did not alter

189 the original lack of preference between the two olfactory stimuli used in the present  
190 tests.

191 Darcin is a male-specific non-volatile urinary protein (MUP20, MW 18893 Da)  
192 previously shown to induce in females a learned olfactory preference for the particular  
193 pattern of urinary volatiles displayed by an individual mouse (Roberts et al., 2010). In  
194 addition, darcin can also induce spatial learning (female mice also remember the  
195 location where it was presented in a test cage, Roberts et al., 2012). Regarding this, we  
196 should keep in mind that spatial learning is also likely to take place in the present  
197 experiments, since we ran a five-minute preference test in which darcin was present  
198 following the first citralva vs. isoamyl acetate test. However, some relevant differences  
199 between the present experiments and those reported by Roberts et al. (2012) suggest a  
200 weaker role of spatial learning in the present case. Firstly, we used a test cage with no  
201 internal spatial cues, with both sides of the cage being equal. Secondly, our test was five  
202 minute long (the training session in Roberts et al., 2012, was 10 minutes long). Thirdly,  
203 we used 8  $\mu$ l of r-darcin, while 50  $\mu$ l were used in Roberts et al. (2012), at  
204 approximately the same concentration. And finally, in the present experiment the  
205 females were later exposed to isoamyl acetate scented papers impregnated with darcin  
206 daily for 15 min over the next four days, with these sessions taking place in a very  
207 different context. This provided abundant possibilities for the formation of odour-  
208 pheromone associations, whereas the opportunities for spatial learning were  
209 comparatively more reduced. In any case, the possible role of spatial learning cannot be  
210 discarded, and future experiments should confirm the induction of odour-pheromone  
211 learning suggested by the present results.

212 Previous work has shown that isoamyl acetate can be used as a conditioned stimulus in  
213 an aversive learning task, associating it with lithium chloride (Kay & Nyby, 1992), and

214 therefore this olfactory stimulus can be conditioned to be either aversive or attractive.

215 We hypothesize that in the present case of associative learning the conditioned stimulus

216 (isoamyl acetate) is detected by the olfactory system and the unconditioned stimulus

217 (darcin) is detected by the vomeronasal system (since the animals need direct contact

218 with the stimulus to show either innate attraction or learned associations, Roberts et al.,

219 2010). Olfactory and vomeronasal information are known to converge in several nuclei

220 within the corticomедial amygdala (Cádiz-Moretti, Martínez-García, & Lanuza, 2013;

221 Kang, Baum, & Cherry, 2009, 2011; Pro-Sistiaga et al., 2007), where learning may take

222 place. In addition, further intramygdaloid projections would allow the participation of

223 the nuclei of the associative (basolateral) amygdala, as suggested by functional data

224 obtained with the immediate early gene c-Fos (Moncho-Bogani, Martínez-García,

225 Novejarque, & Lanuza, 2005). A different pheromone that has been shown to induce

226 olfactory learning is the rabbit mammary pheromone 2-methylbut-2-enal (2MB2)

227 (Coureaud et al., 2006), although in this case the 2MB2 is likely detected by the main

228 olfactory system, as indicated by functional studies using the Fos protein as neural

229 activity marker (Charra, Datiche, Gigot, Schaal, & Coureaud, 2013).

230 The induction of a learned preference for airborne urinary stimuli should take place in

231 natural conditions when female mice explore the urine marks that males use to advertise

232 their territory (Hurst & Beynon, 2004). The urine of males is enriched in several volatile

233 molecules, such as farnesenes, 2-sec-butyl 4,5 dihydrothiazole, and 3,4 dehydro-*exo*-

234 brevicomin, which have been shown to be also detected by the vomeronasal organ

235 (Leinders-Zufall et al., 2000) and maybe possess pheromonal activity on their own (see,

236 for a review, Dulac & Torello, 2003). For example, the mixture of alpha and beta

237 farnesenes was shown to be attractive to sexually naïve female mice but only when used

238 in very high concentrations, while having no effect when used at a concentration that

239 was double that of normal dominant male urine (Jemiolo, Xie, & Novotny, 1991). By  
240 contrast, farnesenes were preferred even at low concentrations by sexually experienced  
241 animals (Jemiolo et al., 1991). Although the previous chemosensory experience of the  
242 animals in these experiments is unknown, the effects of sexual experience clearly  
243 indicate a role for learning. Similar remarks can be made in other cases of putatively  
244 identified pheromonal stimuli, such as (methylthio)methanethiol (MTMT, an attractive  
245 semiochemical present only in the urine of male mice, Lin, Zhang, Block, & Katz,  
246 2005) and androstenone, a pheromone that facilitates expression of both attraction to the  
247 male and a receptive mating stance in estrous female pigs (Dorries, Adkins-Regan, &  
248 Halpern, 1997). In both studies the female subjects had previous sexual experience (in  
249 the case of the female pigs most of them were multiparous). In the light of the results  
250 presented here, the pheromonal role of these semiochemicals should be reevaluated at  
251 least using sexually naïve (if not chemosensory naïve) animals to understand the  
252 requirement for learning. In the same vein, the human steroid androstenone ( $5\alpha$ -androst-  
253 16-en-3-one) has been proposed to function as a human sex chemosignal (see, for a  
254 review, Havlicek, Murray, Saxton, & Roberts, 2010). However, the hedonic value of  
255 androstenone was recently evaluated as a function of sexual experience (Knaapila et al.,  
256 2012), the odour being rated as unpleasant by women who reported never having  
257 experienced sexual intercourse, and as less unpleasant by those who reported being  
258 sexually experienced. Since humans do not have a functional vomeronasal organ  
259 (Meredith, 2001), in this case learning is likely to be mediated by the association of the  
260 olfactory cue with other kinds of rewarding stimuli related to sexual activity.

261 The phenomenon of pheromone-induced olfactory learning raises the question of  
262 whether a substance that gains its role as chemical signal by a learned association  
263 should be considered a pheromone, since it does not elicit a fixed (stereotyped) response

264 (as required by the original definition of Karlson & Luscher, 1953) before learning takes  
265 place. However, in the case of the male-specific airborne urinary substances this  
266 olfactory-vomer nasal association would necessarily occur every time the female  
267 interacts with males of male urine marks. Moreover, females will encounter different  
268 olfactory-vomer nasal associations with each male, so that the learned response would  
269 be specific to that particular signature (Ramm et al., 2008). Under natural conditions,  
270 females are probably able to detect male-specific cues at a distance using the main  
271 olfactory system. Once the female locates the male (or his urine marks) the input  
272 through the vomer nasal organ (requiring direct contact with the source) would allow  
273 further information about this particular individual to be processed. Several of the male-  
274 specific urinary volatiles are also detected by the vomer nasal organ (Leinders-Zufall et  
275 al., 2000), but it is currently unknown why the detection of volatile signals by the  
276 vomer nasal organ requires direct contact with the stimulus, as indicated by both  
277 behavioural and electrophysiological evidence (Luo, Fee, & Katz, 2003; Moncho-  
278 Bogani et al., 2002; Ramm et al., 2008).

279

280 **Turning attraction into aggression: male-specific vomer nasal cues elicit maternal**  
281 **aggression**

282 As stated above, the attraction that females display for male urine is innate. However,  
283 during lactation females show aggressive responses towards male (to a lesser extent also  
284 towards female) intruders, to protect their pups (maternal aggression, Rosenson &  
285 Asheroff, 1975). Maternal aggression towards intruders is observed in the first two  
286 weeks after delivery, and disappears gradually onwards (Lonstein & Gammie, 2002).  
287 Maternal aggression is low towards castrated males, and vomer nasal organ removal in  
288 females prior to mating or after parturition eliminates later maternal aggression in mice

289 (Bean & Wysocki, 1989). Therefore, for the female to show maternal aggression the  
290 vomeronasal detection of testosterone-dependent chemical stimuli from males is  
291 required. This raises the question of whether some of the major urinary proteins, whose  
292 synthesis is testosterone-dependent and have been shown to be the vomeronasal stimuli  
293 mediating aggression between males (Chamero et al., 2007), are also the vomeronasal  
294 stimuli that elicit maternal aggression in lactating females. Preliminary data in our  
295 laboratory suggest that this is indeed the case (Martín-Sánchez et al., 2013), and that the  
296 attractive pheromone darcin is also able to induce maternal aggression. Therefore, the  
297 hormonal status of the lactating females induces changes in the neural structures  
298 processing darcin (probably not in the vomeronasal organ, although direct evidence of  
299 this is lacking). These changes may take place in the amygdalo-hypothalamic circuits  
300 involved in the aggressive response (Nelson & Trainor, 2007), but experimental  
301 evidence of this hypothesis is needed. Notably, lactating females are not aggressive  
302 towards familiar mates (Lonstein & Gammie, 2002), and therefore the response also  
303 depends on the learned identity of the individual male. Individual recognition in mice is  
304 mediated by the pattern of major urinary proteins (Hurst et al., 2001), and therefore the  
305 detection of the pattern of major urinary proteins corresponding to the familiar male  
306 should inhibit the aggressive response. Social recognition involves changes of gene  
307 expression (in particular oxytocin, vasopressin and steroid hormone receptors) in the  
308 amygdala and the hypothalamus (Clipperton-Allen et al., 2012), giving support to the  
309 hypothesis of the control of the aggressive responses by amygdalo-hypothalamic  
310 circuits (Bosch & Neumann, 2012).

311

312 **Turning aversion into attraction: changes at puberty inducing attraction to male**  
313 **pheromones**

314 Although the sexual attraction that female mice display for the male sexual pheromone  
315 darcin does not require learning (Roberts et al, 2010), it seems to appear with puberty,  
316 since pre-pubertal females display an aversive response to chemical signals from  
317 unfamiliar adult males (Drickamer, 1989; Mucignat-Caretta, Caretta, & Baldini, 1998).  
318 The biological bases of this change are unknown, although clearly the gonadal steroids  
319 underlying the pubertal changes should play a relevant role. In fact, the vomeronasal  
320 organ, likely involved in the detection of protein sex pheromones (such as darcin,  
321 Roberts et al., 2010; and exocrine gland-secreting peptide 1, Haga et al., 2010), as well  
322 as most of the neural centres of the vomeronasal system are sexually dimorphic  
323 (Segovia & Guillaumon, 1993). This suggests a role for sexual steroids in development  
324 and differentiation of the vomeronasal system. Moreover, gonadal steroids also regulate  
325 the expression of some vomeronasal receptors (Alekseyenko, Baum, & Cherry, 2006),  
326 raising the possibility that sex steroids induce the expression of particular (currently  
327 unknown) receptors for male pheromones. In fact, there is some evidence of estradiol  
328 effects on the induction of c-Fos in the vomeronasal organ of female mice by male-  
329 soiled bedding (Halem, Cherry, & Baum, 1999). In addition, the presence of estradiol  
330 receptors is very important in the secondary vomeronasal centres, namely the medial  
331 amygdala, the posteromedial cortical amygdala and the posteromedial bed nucleus of  
332 the stria terminalis (Mitra et al., 2003). Therefore, the pubertal changes underlying the  
333 induction of the attraction of female mice for male pheromones may also take place in  
334 the amygdaloid circuits processing vomeronasal information.

335

336 **From attraction to aversion: illness-derived chemicals and avoidance of**  
337 **conspecifics**

338 During the last years, our research has been focused on the attractive properties of  
339 sexual pheromones and neurobiological foundation of this phenomenon. Occasionally  
340 we observed that viral infections or parasitosis in male mice used as donors of urine or  
341 bedding resulted in a lack of attractiveness of the urine (or soiled bedding) for females.  
342 This fits previous studies reporting avoidance of males by females when males were  
343 infected with viruses or parasites (e.g. Penn et al. 1998; Kavaliers et al., 2005). To  
344 check the response of female mice to the urine of infected males experimentally, we  
345 performed preference tests (Experiment 2) using urine from males parasitized by the  
346 nematode *Aspiculuris tetraptera* against (healthy) female urine. In addition, since darcin  
347 appears to be the both necessary and sufficient to make male urine attractive to females,  
348 we tested whether the urine of parasitized males contains darcin.

349 EXPERIMENT 2. Behavioural response of female mice to the urine of infected males

350 Material and Methods

351 *Subjects and stimuli*

352 For this experiment, 32 adult female mice (12-16 weeks) of CD1 strain were used  
353 (Janvier, Le Genest-Saint-Isle, Saint-Berthevin Cedex, France). As for experiment 1,  
354 females were sexually naïve and had never been exposed to chemical signals from  
355 sexually mature males. Females were housed in groups of 5-6 animals, with the housing  
356 conditions (cages, bedding and food), manipulation and welfare assessment being the  
357 same as those reported in Experiment 1.

358 Urine from healthy adult CD-1 male and female mice was purchased from Janvier and  
359 kept frozen in aliquots until used. Urine from a small colony (n = 4) of male mice (also  
360 purchased from Janvier) naturally infected with the intestinal nematode *Aspiculuris*  
361 *tetraptera* was collected as described by Kurien, Everds & Scofield (2004). Briefly,  
362 animals were gently held by the scruff of the neck over a petri dish, from where urine



363 was pipetted. Since the infection occurred naturally, the animals can probably  
364 experience this level of parasitism in the wild. The presence of these parasites in male  
365 mice was detected with the routine sentinel vigilance system. The infected mice were  
366 sacrificed except for four animals that were kept for 5 days. During this time we  
367 collected their urine once a day. At the end of this 5-day period the four infected  
368 animals were also sacrificed. The infected mice showed no external signs of infection,  
369 and general appearance, size, coat condition, posture, gait, activity levels, interaction  
370 with the environment and clinical signs appeared normal. Infected male mice were  
371 housed in pairs in standard macrolon cages with a wire lid (22.5 x 22.5 x 14.5 cm, ref.  
372 500, Panlab, Barcelona, Spain). The rest of housing conditions were the same as  
373 described in Experiment 1. To ensure the homogeneity of the stimulus across  
374 behavioural tests, urine from different males was mixed and stored in frozen aliquots of  
375 50 µl. Infection with the parasite was assessed through the presence of eggs in faecal  
376 pellets and confirmed post-mortem by checking for adult worms in the colon.

### 377 *Preference tests*

378 The test cage and habituation procedure were as described in experiment 1. Preference  
379 tests were performed in which the female mice had to choose between two urine stimuli  
380 located in opposite sides of the cage. To do so, 10 µl of the stimulus urine were pipetted  
381 on one of the tips of a rectangular piece of filter paper (2 x 6 cm) that was attached to  
382 the wall so that the urine spots were 8 cm above the floor. The females were able to  
383 have direct nasal contact with the stimuli by standing on the hind legs. Following a  
384 control test, with PBS 0.1M on both sides of the cage, the olfactory preference test  
385 (male vs female urine) was performed and recorded for five minutes as described for  
386 Experiment 1. In one group of animals (randomly assigned, n =16), urine of healthy  
387 males was presented on one side and female urine on the other side. In the second group

388 of females (randomly assigned, n = 16), urine of infected males was presented against  
389 female urine.

#### 390 *Statistical analysis*

391 The time spent in a semicircular area (of a radius of 3.85 cm) around the filter paper was  
392 analysed with a two-way repeated-measures ANOVA with TEST (control, preference  
393 test) and STIMULUS (male vs. female urine) as intra-subject factors, and GROUP  
394 (urine from healthy or infected males) as an inter-subject factor. Significant interactions  
395 were further analysed by multiple pair-wise comparisons with Bonferroni corrections.  
396 The normality of the data was previously confirmed with a Kolmogorov-Smirnov test  
397 with Lillieford's correction. Analyses were performed with the SPSS 15.0 software  
398 package.

#### 399 *Electrophoresis of urinary proteins*

400 The pattern of bands corresponding to urinary proteins, separated according to their  
401 mass, was visualised using sodium dodecyl sulphate–polyacrylamide gel electrophoresis  
402 (SDS–PAGE). Urine was diluted 1:1 with 2x denaturalization buffer (20mM Tris pH  
403 8.0, 5% SDS, 10% mercaptoethanol, 2mM EDTA and 0.05% bromophenol blue) in a  
404 capped Eppendorf tube, vortexed to mix, boiled for 5 min and then centrifuged for 5  
405 min at 10000 rpm. The samples were then allowed to cool before sample loading. Using  
406 1x denaturalisation buffer samples of male urine were brought to a final 1:6 dilution,  
407 whereas female urine samples were not further diluted (final dilution 1:2). This allowed  
408 direct comparison of male and female urine protein species, in spite of the difference in  
409 total urinary protein content between sexes. Using a PhastGel system (General  
410 Electrics), electrophoresis was run under reducing conditions at a constant 200 V on a  
411 20% polyacrylamide gel (PhastGel Homogeneous – 20, GE). Low range molecular  
412 weight markers (Sigmamarker low range, M3913, St. Louis, MO, USA) were used for

413 comparison. Following electrophoresis, protein bands were visualised using Phast Gel  
414 Blue (0.1%) solution and differentiated in a solution of methanol: acetic acid: distilled  
415 H<sub>2</sub>O (30:10:60 v/v/v).

## 416 Results

417 The results of the repeated measures ANOVA of time spent within the areas  
418 surrounding the stimulus during the tests (Figure 2A) showed non-significant main  
419 effects of factors TEST ( $F_{1,30} = 1.9$ ,  $p = 0.17$ ), STIMULUS ( $F_{1,30} < 1$ ,  $p = 0.6$ ) or  
420 GROUP ( $F_{1,30} < 1$ ,  $p = 0.9$ ), and non-significant interactions between each pair of these  
421 factors (STIMULUS x TEST:  $F_{1,30} = 1.14$ ,  $p = 0.29$ ; TEST x GROUP:  $F_{1,30} = 2.43$ ,  $p =$   
422  $0.12$ ; STIMULUS x GROUP:  $F_{1,30} < 1$ ,  $p = 0.4$ ). However, there was a significant triple  
423 interaction (STIMULUS x TEST x GROUP:  $F_{1,30} = 7.45$ ,  $p = 0.01$ ). Pos-hoc analysis of  
424 this triple interaction (Figure 2A) showed that in both groups the two stimulus areas  
425 were investigated equally in the control condition, when saline buffer was present on  
426 both sides of the cage (healthy male group,  $p > 0.7$ ; infected male group,  $p > 0.9$ ). By  
427 contrast, females presented with urine of healthy males showed a clear preference for  
428 this stimulus over female urine ( $p = 0.023$ ); females presented with urine of infected  
429 males spent more time next to the female urine, although the time spent next to each  
430 stimulus was not significantly different ( $p = 0.27$ ).

431 To check for the presence of the sexual pheromone darcin in the urine of male mice  
432 infected with the nematode *Aspicularis tetraptera*, we performed a SDS-PAGE  
433 electrophoresis to compare proteins in healthy male urine, urine of infected males, urine  
434 of castrated (non-infected) males and female urine (Figure 2B). The results showed that  
435 the pattern of protein bands in the urine of infected males was similar to that observed  
436 in the urine of healthy males, with a clearly visible band of higher mobility than other  
437 major urinary proteins that corresponds to darcin (Armstrong et al., 2005). By contrast,

438 this band was not present in the urine from castrated males or in female urine,  
439 confirming previous results (Armstrong, Robertson, Cheetham, Hurst, & Beynon, 2005;  
440 Cheetham, Smith, Armstrong, Beynon, & Hurst, 2009).

#### 441 Discussion

442 It has been shown previously that female mice are able to discriminate between infected  
443 and non-infected males and show a preference towards healthy males (see Kavaliers,  
444 Choleris, & Pfaff, 2005). The results of the present experiment suggest that females not  
445 only choose non-infected males against infected males, but also that the preference for  
446 male urine over female urine is lost if urine comes from parasitized males. Since it is  
447 known that the attractive properties of male urine depend on the presence of a male-  
448 specific urinary protein named darcin (Roberts et al., 2010), which acts as a sexual  
449 pheromone, we tested whether the expression of this protein may have been lost in the  
450 infected males. The results show that darcin is present in the urine of parasitized males  
451 apparently at normal levels, though further analyses would be required to confirm  
452 whether there are quantitative differences. Although we cannot discard a small  
453 reduction in the expression of this protein, this is unlikely to explain the total lack of  
454 preference for male urine versus female urine that did not contain darcin (at least at a  
455 level that could be detected by electrophoresis). Changes in the amino-acidic sequence  
456 or conformation of the protein are very unlikely. Therefore, we can hypothesize that an  
457 infection cue exists in the urine of the parasitized animals, and that detection of this is  
458 able to override the attractive value of darcin. An alternative hypothesis, though, is that  
459 the cue of infection is volatile and can be detected at a distance; this detection may  
460 inhibit the vomeronasal pumping necessary to deliver high molecular weight molecules,  
461 such darcin, to the vomeronasal organ (Meredith, Marques, O'Connell, & Stern, 1980;

462 Wysocki, Wellington, & Beauchamp, 1980). In this case, darcin would simply not be  
463 detected by females.

464 Whether volatile or involatile (or both), the identity of the putative cue(s) of infection is  
465 unknown, as it is its nature as olfactory of vomeronasal stimulus. It has recently been  
466 demonstrated that the vomeronasal organ of mice expresses a different type of  
467 chemosensory receptor named FPR (formyl peptide receptors, Liberles et al., 2009;  
468 Riviere, Challet, Fluegge, Spehr, & Rodriguez, 2009). This type of receptor, in addition  
469 to formylated peptides produced by bacteria, detects ligands related with the immune  
470 system (such the antimicrobial peptide CRAMP, lipoxin A4, or uPAR, see Riviere et  
471 al., 2009), and therefore are good candidates for sensing of urinary infection cues,  
472 although experimental evidence for this possibility is currently lacking.

473

#### 474 GENERAL CONCLUSIONS

475 Although the attractive response that adult female mice show towards the male-specific  
476 pheromone darcin is innate, here we review four examples in which either the  
477 behavioural response can be induced by non-pheromonal stimuli or the presence of the  
478 pheromone does not induce the normal attractive response, due a non-receptive  
479 hormonal status (pre-pubertal or lactating females) or to the putative presence of  
480 infection cues in the male urine.

481 The induction of learned attraction by a pheromone is an interesting model to study  
482 learning and memory mechanisms in an ethologically relevant context. In addition,  
483 changes in response to darcin induced at puberty and during lactation provide  
484 unexplored models of neural plasticity with clear and robust behavioural correlates,  
485 which may be helpful to understand the neural circuits that induce both attractive and  
486 aggressive responses.

487

488 **ACKNOWLEDGEMENTS**

489 Funded by the Spanish Ministry of Science-FEDER (BFU2010-16656) and the UK  
490 Biotechnology and Biological Sciences Research Council (BB/J002631/1). The authors  
491 are grateful to Dr. Antonio Marcilla and Dr. Maria Trelis (Dept. of Parasitology,  
492 University of Valencia) for their invaluable help to deal with the infected mice.

493 REFERENCES

- 494 **Alekseyenko, O. V., Baum, M. J. & Cherry, J. A.** 2006. Sex and gonadal steroid  
495 modulation of pheromone receptor gene expression in the mouse vomeronasal organ.  
496 *Neuroscience*, **140**, 1349-1357. doi: 10.1016/j.neuroscience.2006.03.001.
- 497 **Angely, C. J. & Coppola, D. M.** 2010. How does long-term odor deprivation affect the  
498 olfactory capacity of adult mice? *Behavioral and Brain Functions*, **6**, 26-9081-6-26.  
499 doi: 10.1186/1744-9081-6-26.
- 500 **Armstrong, S. D., Robertson, D. H., Cheetham, S. A., Hurst, J. L. & Beynon, R. J.**  
501 2005. Structural and functional differences in isoforms of mouse major urinary proteins:  
502 a male-specific protein that preferentially binds a male pheromone. *The Biochemical*  
503 *Journal*, **391**, 343-350.
- 504 **Bean, N. J. & Wysocki, C. J.** 1989. Vomeronasal organ removal and female mouse  
505 aggression: the role of experience. *Physiology & Behavior*, **45**, 875-882.
- 506 **Bosch, O.J. & Neumann, I.D.** 2012. Both oxytocin and vasopressin are mediators of  
507 maternal care and aggression in rodents: from central release to sites of action.  
508 *Hormones and Behavior*, **61**, 293-303. doi: 10.1016/j.yhbeh.2011.11.002.
- 509 **Brennan, P. A. & Kendrick, K. M.** 2006. Mammalian social odours: attraction and  
510 individual recognition. *Philosophical Transactions of the Royal Society. Series B,*  
511 *Biological Sciences*, **361**, 2061-2078. doi: 10.1098/rstb.2006.1931.
- 512 **Cadiz-Moretti, B., Martinez, G. F. & Lanuza, E.** 2013. Neural substrate to associate  
513 odorants and pheromones: Convergence of projections from the main and accessory  
514 olfactory bulbs in mice. In: *Chemical Signals in Vertebrates 12* (Ed. by M. L. East & M.  
515 Dehnhard), pp. 3-16. New York: Springer Science.
- 516 **Chamero, P., Marton, T. F., Logan, D. W., Flanagan, K., Cruz, J. R., Saghatelian,**  
517 **A., Cravatt, B. F. & Stowers, L.** 2007. Identification of protein pheromones that  
518 promote aggressive behaviour. *Nature*, **450**, 899-902. doi: 10.1038/nature05997.

- 519 **Charra, R., Datiche, F., Gigot, V., Schaal, B., & Coureaud, G.** 2013. Pheromone-  
520 induced odor learning modifies Fos expression in the newborn rabbit brain. *Behavioral*  
521 *Brain Research*, **237**, 129-140. doi: 10.1016/j.bbr.2012.09.017.
- 522 **Cheetham, S. A., Smith, A. L., Armstrong, S. D., Beynon, R. J. & Hurst, J. L.** 2009.  
523 Limited variation in the major urinary proteins of laboratory mice. *Physiology &*  
524 *Behavior*, **96**, 253-261. doi: 10.1016/j.physbeh.2008.10.005.
- 525 **Clipperton-Allen, A.E., Lee, A.W., Reyes, A., Devidze, N., Phan, A., Pfaff, D.W. &**  
526 **Choleris, E.** 2012. Oxytocin, vasopressin and estrogen receptor gene expression in  
527 relation to social recognition in female mice. *Physiology and Behavior*, **105**, 915-924.  
528 doi: 10.1016/j.physbeh.2011.10.025.
- 529 **Coureaud, G., Moncomble, A.S., Montigny, D., Dewas, M., Perrier, G., & Schaal,**  
530 **B.** 2006. A pheromone that rapidly promotes learning in the newborn. *Current Biology*,  
531 **16**, 1956-1961.
- 532 **Dorries, K. M., Adkins-Regan, E. & Halpern, B. P.** 1997. Sensitivity and behavioral  
533 responses to the pheromone androstenone are not mediated by the vomeronasal organ in  
534 domestic pigs. *Brain, Behavior and Evolution*, **49**, 53-62.
- 535 **Drickamer, L. C.** 1989. Odor preference of wild stock female house mice (*Mus*  
536 *domesticus*) tested at three ages using urine and other cues from conspecific males and  
537 females. *Journal of Chemical Ecology*, **15**, 1971-1987.
- 538 **Dulac, C. & Torello, A. T.** 2003. Molecular detection of pheromone signals in  
539 mammals: from genes to behaviour. *Nature Reviews Neuroscience*, **4**, 551-562.
- 540 **Haga, S., Hattori, T., Sato, T., Sato, K., Matsuda, S., Kobayakawa, R., Sakano, H.,**  
541 **Yoshihara, Y., Kikusui, T. & Touhara, K.** 2010. The male mouse pheromone ESP1  
542 enhances female sexual receptive behaviour through a specific vomeronasal receptor.  
543 *Nature*, **466**, 118-122. doi: 10.1038/nature09142.
- 544 **Halem, H. A., Cherry, J. A. & Baum, M. J.** 1999. Vomeronasal neuroepithelium and  
545 forebrain Fos responses to male pheromones in male and female mice. *Journal of*  
546 *Neurobiology*, **39**, 249-263.



- 547 **Havlicek, J., Murray, A. K., Saxton, T. K. & Roberts, S. C.** 2010. Current issues in  
548 the study of androstenes in human chemosignaling. *Vitamins and Hormones*, **83**, 47-81.  
549 doi: 10.1016/S0083-6729(10)83003-1.
- 550 **Herz, R. S. & Cupchik, G. C.** 1995. The emotional distinctiveness of odor-evoked  
551 memories. *Chemical Senses*, **20**, 517-528.
- 552 **Honda, K., Negoro, H., Dyball, R. E. J., Higuchi, T. & Takano, S.** 1990. The  
553 osmoreceptor complex in the rat: Evidence for interactions between the supraoptic and  
554 other diencephalic nuclei. *The Journal of Physiology*, **431**, 225-241.
- 555 **Hurst, J.L., Payne, C.E., Nevison, C.M., Marie, A.D., Humphries, R.E., Robertson,**  
556 **D.H., Cavaggioni, A. & Beynon, R.J.** 2001. Individual recognition in mice mediated  
557 by major urinary proteins. *Nature*, **414**, 631-634.
- 558 **Hurst, J. L. & Beynon, R. J.** 2004. Scent wars: the chemobiology of competitive  
559 signalling in mice. *BioEssays*, **26**, 1288-1298.
- 560 **Jemiolo, B., Xie, T. M. & Novotny, M.** 1991. Socio-sexual olfactory preference in  
561 female mice: attractiveness of synthetic chemosignals. *Physiology & Behavior*, **50**,  
562 1119-1122.
- 563 **Kang, N., Baum, M. J. & Cherry, J. A.** 2011. Different profiles of main and accessory  
564 olfactory bulb mitral/tufted cell projections revealed in mice using an anterograde tracer  
565 and a whole-mount, flattened cortex preparation. *Chemical Senses*, **36**, 251-260. doi:  
566 10.1093/chemse/bjq120; 10.1093/chemse/bjq120.
- 567 **Kang, N., Baum, M. J. & Cherry, J. A.** 2009. A direct main olfactory bulb projection  
568 to the 'vomeronasal' amygdala in female mice selectively responds to volatile  
569 pheromones from males. *The European Journal of Neuroscience*, **29**, 624-634. doi:  
570 10.1111/j.1460-9568.2009.06638.x.
- 571 **Karlson, P. & Luscher, M.** 1959. Pheromones': a new term for a class of biologically  
572 active substances. *Nature*, **183**, 55-56.

- 573 **Kavaliers, M., Choleris, E. & Pfaff, D. W.** 2005. Genes, odours and the recognition of  
574 parasitized individuals by rodents. *Trends in Parasitology*, **21**, 423-429. doi:  
575 10.1016/j.pt.2005.07.008.
- 576 **Kay, E. & Nyby, J.** 1992. LiCl aversive conditioning has transitory effects on  
577 pheromonal responsiveness in male house mice (*Mus domesticus*). *Physiology &*  
578 *Behavior*, **52**, 105-113.
- 579 **Knaapila, A., Tuorila, H., Vuoksima, E., Keskitalo-Vuokko, K., Rose, R. J.,**  
580 **Kaprio, J. & Silventoinen, K.** 2012. Pleasantness of the odor of androstenone as a  
581 function of sexual intercourse experience in women and men. *Archives of Sexual*  
582 *Behavior*, **41**, 1403-1408. doi: 10.1007/s10508-011-9804-7.
- 583 **Kurien, B.T., Everds, N.E. & Scofield, R.H.** 2004. Experimental animal urine  
584 collection: a review. *Laboratory Animals*, **38**, 333-361.
- 585 **Leinders-Zufall, T., Lane, A. P., Puche, A. C., Ma, W., Novotny, M. V., Shipley, M.**  
586 **T. & Zufall, F.** 2000. Ultrasensitive pheromone detection by mammalian vomeronasal  
587 neurons. *Nature*, **405**, 792-796.
- 588 **Liberles, S. D., Horowitz, L. F., Kuang, D., Contos, J. J., Wilson, K. L., Siltberg-**  
589 **Liberles, J., Liberles, D. A. & Buck, L. B.** 2009. Formyl peptide receptors are  
590 candidate chemosensory receptors in the vomeronasal organ. *Proceedings of the*  
591 *National Academy of Sciences USA*, **106**, 9842-9847. doi: 10.1073/pnas.0904464106.
- 592 **Lonstein, J. S. & Gammie, S. C.** 2002. Sensory, hormonal, and neural control of  
593 maternal aggression in laboratory rodents. *Neuroscience and Biobehavioral Reviews*,  
594 **26**, 869-888.
- 595 **Luo, M., Fee, M. S. & Katz, L. C.** 2003. Encoding pheromonal signals in the  
596 accessory olfactory bulb of behaving mice. *Science*, **299**, 1196-1201. doi:  
597 10.1126/science.1082133.
- 598 **Martin-Sanchez, A., Hernandez-Martinez, A., McLean, L., Beynon, R.J., Hurst,**  
599 **J.L., Lanuza, E., Martinez-Garcia, F.** 2013. When males become the enemy: maternal

600 aggression is induced by the attractive male sexual pheromone darcin in mice. 5th  
601 Parental Brain Conference. Póster. Regensburg, Germany.

602 **Martinez-Garcia, F., Martinez-Ricos, J., Agustin-Pavon, C., Martinez-Hernandez,**  
603 **J., Novejarque, A. & Lanuza, E.** 2009. Refining the dual olfactory hypothesis:  
604 pheromone reward and odour experience. *Behavioural Brain Research*, **200**, 277-286.  
605 doi: 10.1016/j.bbr.2008.10.002.

606 **Martinez-Ricos, J., Agustin-Pavon, C., Lanuza, E. & Martinez-Garcia, F.** 2007.  
607 Intraspecific communication through chemical signals in female mice: reinforcing  
608 properties of involatile male sexual pheromones. *Chemical Senses*, **32**, 139-148.

609 **Martinez-Ricos, J., Agustin-Pavon, C., Lanuza, E. & Martinez-Garcia, F.** 2008.  
610 Role of the vomeronasal system in intersexual attraction in female mice. *Neuroscience*,  
611 **153**, 383-395. doi: 10.1016/j.neuroscience.2008.02.002.

612 **Meredith, M.** 2001. Human vomeronasal organ function: a critical review of best and  
613 worst cases. *Chemical Senses*, **26**, 433-445.

614 **Meredith, M., Marques, D. M., O'Connell, R. O. & Stern, F. L.** 1980. Vomeronasal  
615 pump: significance for male hamster sexual behavior. *Science*, **207**, 1224-1226.

616 **Mitra, S. W., Hoskin, E., Yudkovitz, J., Pear, L., Wilkinson, H. A., Hayashi, S.,**  
617 **Pfaff, D. W., Ogawa, S., Rohrer, S. P., Schaeffer, J. M., McEwen, B. S. & Alves, S.**  
618 **E.** 2003. Immunolocalization of estrogen receptor beta in the mouse brain: comparison  
619 with estrogen receptor alpha. *Endocrinology*, **144**, 2055-2067.

620 **Moncho-Bogani, J., Martinez-Garcia, F., Novejarque, A. & Lanuza, E.** 2005.  
621 Attraction to sexual pheromones and associated odorants in female mice involves  
622 activation of the reward system and basolateral amygdala. *The European Journal of*  
623 *Neuroscience*, **21**, 2186-2198.

624 **Moncho-Bogani, J., Lanuza, E., Hernandez, A., Novejarque, A. & Martinez-**  
625 **Garcia, F.** 2002. Attractive properties of sexual pheromones in mice. Innate or learned?  
626 *Physiology & Behavior*, **77**, 167-176.

- 627 **Mucignat-Caretta, C., Caretta, A. & Baldini, E.** 1998. Protein-bound male urinary  
628 pheromones: differential responses according to age and gender. *Chemical Senses*, **23**,  
629 67-70.
- 630 **Nelson, R. J. & Trainor, B. C.** 2007. Neural mechanisms of aggression. *Nature*  
631 *Reviews Neuroscience*, **8**, 536-546. doi: 10.1038/nrn2174.
- 632 **Otero-Garcia, M., Martín-Sánchez, A., Fortes-Marco, L., Martínez-Ricós, J.,**  
633 **Agustín-Pavón, C., Lanuza, E. & Martínez-García, F.** 2014. Extending the socio-  
634 sexual brain: arginine-vasopressin immunoreactive circuits in the telencephalon of mice.  
635 *Brain Structure and Function* (in press). Doi: 10.1007/s00429-013-0553-3.
- 636 **Penn, D., Schneider, G., White, K., Slev, P. & Potts, W.** 2010. Influenza infection  
637 neutralizes the attractiveness of male odour to female mice (*Mus musculus*). *Ethology*,  
638 **104**, 685-694. doi: 10.1111/j.1439-0310.1998.tb00102.x.
- 639 **Pro-Sistiaga, P., Mohedano-Moriano, A., Ubeda-Banon, I., Del Mar Arroyo-**  
640 **Jimenez, M., Marcos, P., Artacho-Perula, E., Crespo, C., Insausti, R. & Martinez-**  
641 **Marcos, A.** 2007. Convergence of olfactory and vomeronasal projections in the rat  
642 basal telencephalon. *The Journal of Comparative Neurology*, **504**, 346-362. doi:  
643 10.1002/cne.21455.
- 644 **Ramm, S. A., Cheetham, S. A. & Hurst, J. L.** 2008. Encoding choosiness: female  
645 attraction requires prior physical contact with individual male scents in mice.  
646 *Proceedings of the Royal Society, Series B, Biological Sciences*, **275**, 1727-1735. doi:  
647 10.1098/rspb.2008.0302; 10.1098/rspb.2008.0302.
- 648 **Riviere, S., Challet, L., Fluegge, D., Spehr, M. & Rodriguez, I.** 2009. Formyl peptide  
649 receptor-like proteins are a novel family of vomeronasal chemosensors. *Nature*, **459**,  
650 574-577. doi: 10.1038/nature08029.
- 651 **Roberts, S. A., Simpson, D. M., Armstrong, S. D., Davidson, A. J., Robertson, D.**  
652 **H., McLean, L., Beynon, R. J. & Hurst, J. L.** 2010. Darcin: a male pheromone that  
653 stimulates female memory and sexual attraction to an individual male's odour. *BMC*  
654 *Biology*, **8**, 75-7007-8-75. doi: 10.1186/1741-7007-8-75.

655 **Roberts, S. A., Davidson, A. J., McLean, L., Beynon, R. J. & Hurst, J. L.** 2012.  
656 Pheromonal induction of spatial learning in mice. *Science*, **338**, 1462-1465. doi:  
657 10.1126/science.1225638.

658 **Rosenson, L. M. & Asheroff, A. K.** 1975. Maternal aggression in CD-1 mice:  
659 Influence of the hormonal condition of the intruder. *Behavioral Biology*, **15**, 219-224.  
660 doi: 10.1016/S0091-6773(75)91603-X.

661 **Segovia, S. & Guillamon, A.** 1993. Sexual dimorphism in the vomeronasal pathway  
662 and sex differences in reproductive behaviors. *Brain Research Reviews*, **18**, 51-74.

663 **Wells, D.J., Playle, L.C., Enser, W.E., Flecknell, P.A., Gardiner, M.A., Holland, J.,**  
664 **Howard, B.R., Hubrecht, R., Humphreys, K.R., Jackson, I.J., Lane, N.,**  
665 **Maconochie, M., Mason, G., Morton, D.B., Raymond, R., Robinson, V., Smith,**  
666 **J.A., Watt, N.** 2006. Assessing the welfare of genetically altered mice. *Laboratory*  
667 *Animals* 40:111-114.

668 **Wyatt, T. D.** 2010. Pheromones and signature mixtures: defining species-wide signals  
669 and variable cues for identity in both invertebrates and vertebrates. *Journal of*  
670 *Comparative Physiology A*, **196**, 685-700. doi: 10.1007/s00359-010-0564-y.

671 **Wyatt, T. D.** 2009. Fifty years of pheromones. *Nature*, **457**, 262-263. doi:  
672 10.1038/457262a.

673 **Wysocki, C. J., Wellington, J. L. & Beauchamp, G. K.** 1980. Access of urinary  
674 nonvolatiles to the mammalian vomeronasal organ. *Science*, **207**, 781-783.

675

676

677        FIGURE LEGENDS

678        Figure 1. **A:** Example videotrack of the exploratory behaviour of one animal in an  
679        odour preference test (citralsa versus isoamyl acetate). The white pieces of scented  
680        paper are visible inside the areas of measure. **B:** Time (mean  $\pm$  SE) spent by female  
681        mice in the areas where the odorant stimuli were presented, in the experimental (left  
682        panel, darcin group) and control (right panel) groups. Grey bars in the control test  
683        represent clean pieces of paper. Orange bars represent time spent in the citralsa area.  
684        Green bars represent time spent in the isoamyl acetate area. In test 2 in the experimental  
685        group, the isoamyl acetate-scented paper was also impregnated with r-darcin. Females  
686        showed a significant innate preference to remain near to the r-darcin sample ( $p = 0.032$ )  
687        and a learned preference to stay next to the isoamyl acetate-scented paper in the post  
688        training preference test ( $p = 0.031$ ). In the absence of darcin, no preference appeared for  
689        any odorant.


690

691        Figure 2. **A:** Time (mean  $\pm$  SE) spent by female mice in areas where urine stimuli  
692        were presented. The left side corresponds to females that were presented with urine of  
693        healthy males versus females. Grey bars represent time spent near pieces of paper with a  
694        sample of PBS (control). The green bar represents time spent near urine of healthy  
695        males, and purple bar the time spent near female urine. The right side corresponds to  
696        females that were presented with urine of parasitized males versus females. Grey bars  
697        represent the control test. Females showed no preference for the urine of infected males  
698        (dark red bar) versus female urine (purple bar). **B:** SDS-PAGE of urinary protein for  
699        healthy male mice ( $n = 2$ ), male mice parasitized with *Aspiculuris tetraptera* ( $n = 2$ ),  
700        castrated male mice ( $n = 1$ ) and females ( $n = 2$ ). The 20 kDa band present in all cases  
701        corresponds to the molecular weight to the major urinary proteins. The small band with

702 higher mobility (around 17 kDa), present in healthy and infected males, corresponds to  
703 the expected position of darcin.

Figure 1

**A**



**B**






Figure 2

