

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/341882889>

Health literacy: A crucial determinant of vaccination decision-making

Article in *International Journal of Infectious Diseases* · June 2020

DOI: 10.1016/j.ijid.2020.05.081

CITATIONS

0

READS

16

4 authors:



Enrique Castro-Sánchez

City, University of London

167 PUBLICATIONS 1,128 CITATIONS

[SEE PROFILE](#)



Desirée Mena-Tudela

Universitat Jaume I

56 PUBLICATIONS 50 CITATIONS

[SEE PROFILE](#)



Francisco Javier Soriano-Vidal

Agència Valenciana de Salut

24 PUBLICATIONS 56 CITATIONS

[SEE PROFILE](#)



Rafael Vila-Candel

Agència Valenciana de Salut

70 PUBLICATIONS 114 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Health literacy in pregnancy [View project](#)



Care prescription according to the patient's functional capacity [View project](#)



Letter to the Editor

Health literacy: A crucial determinant of vaccination decision-making

Dear Editor,

We read with interest the Review by [Buchy et al. \(2020\)](#) summarizing the evidence on the micro (patients, healthcare workers), meso (vaccine manufacturers), and macro (wider society) determinants of influenza vaccination in pregnant women worldwide.

We agree with the authors that improvements in vaccination coverage could be gained with better training for healthcare workers about communicating the benefits of vaccination and the risks associated with influenza infection, as well as education about such benefits to pregnant women. However, education per se may be insufficient unless it is underpinned by health literacy principles. Health literacy has been defined primarily as the capacity that citizens have to access and make use of health information to make decisions about their health and care ([Nutbeam, 2008](#)).

Decisions to vaccinate are complex, requiring an understanding of the immune system and the pathogenesis of each microorganism, the variety of immunization schedules, the adverse events which occur at different rates and the influence of several health professionals involved in the decision-making process at different time points ([Vila-Candel et al., 2016a](#)).

Our study exploring the influence of health literacy on influenza vaccination among pregnant women in Spain reflected how women with high health literacy were more likely to reject influenza vaccination than those with lower health literacy ([Castro-Sánchez et al., 2018](#)). Among the reasons for these women to reject the vaccine were feelings about its need, in isolation and compared with other immunizations, and claims of having received insufficient information. To mitigate such a gap, women with high health literacy would access a greater variety of information sources, including those promoting outlier or fringe opinions ([Danchin et al., 2018](#); [Vila-Candel et al., 2016b](#)).

The implications of this study are clear – a full understanding by healthcare professionals of how pregnant women make decisions about vaccinations is required, including the crucial topics of women's interest and concern, which information sources they prefer to meet their needs, how they combine and integrate the evidence obtained into explanatory frameworks, and when these frameworks are shaped by emotions and influenced by social and personal networks.

Interventions solely interested in providing evidence-focused vaccination messages to mitigate assumed information gaps, and without acting upon the determinants shared by vaccination and health literacy such as education level, self-efficacy, and empowerment, are unlikely to sustainably dent existing suboptimal vaccination rates among pregnant women, precluding them from the clinical, public health and social gains afforded by vaccination.

Funding

Enrique Castro-Sánchez is affiliated with the National Institute for Health Research (NIHR) Health Protection Research Unit in Healthcare Associated Infections and Antimicrobial Resistance at Imperial College London (grant No. HPRU- 2012e10047) in collaboration with Public Health England and Imperial College Healthcare NHS Trust. He is also an NIHR Senior Nurse and Midwife Research Leader and recognizes the support of the NIHR Imperial Patient Safety Translational Research Centre and the BRC.

References

- Buchy P, Badur S, Kassianos G, Preiss S, Tam JS. Vaccinating pregnant women against influenza needs to be a priority for all countries: an expert commentary. *Int J Infect Dis* 2020;92:1–12, doi:<http://dx.doi.org/10.1016/j.ijid.2019.12.019>.
- Castro-Sánchez E, Vila-Candel R, Soriano-Vidal FJ, Navarro-Illana E, Díez-Domingo J. Influence of health literacy on acceptance of influenza and pertussis vaccinations: a cross-sectional study among Spanish pregnant women. *BMJ Open* 2018;8(7):e022132, doi:<http://dx.doi.org/10.1136/bmjopen-2018-022132>.
- Danchin MH, Costa-Pinto J, Attwell K, Willaby H, Wiley K, Hoq M, et al. Vaccine decision-making begins in pregnancy: correlation between vaccine concerns, intentions and maternal vaccination with subsequent childhood vaccine uptake. *Vaccine* 2018;36(44):6473–9, doi:<http://dx.doi.org/10.1016/j.vaccine.2017.08.003>.
- Nutbeam D. The evolving concept of health literacy. *Soc Sci Med* 2008;67:2072–8.
- Vila-Candel R, Soriano-Vidal FJ, Castro-Sánchez E. Midwives, vaccines and health literacy: an underexplored relationship?. *Gac Sanit* 2016a;30(5):403, doi:<http://dx.doi.org/10.1016/j.gaceta.2016.05.004>.
- Vila-Candel R, Navarro-Illana P, Navarro-Illana E, Castro-Sánchez E, Duke K, Soriano-Vidal FJ, et al. Determinants of seasonal influenza vaccination in pregnant women in Valencia, Spain. *BMC Public Health* 2016b;16(1):1173, doi:<http://dx.doi.org/10.1186/s12889-016-3823-1>.

Enrique Castro-Sánchez*

NIHR Health Protection Research Unit in Healthcare-Associated Infection and Antimicrobial Resistance at Imperial College London, Du Cane Road, London, W12 0NN, UK

Desirée Mena-Tudela

Department of Nursing, Faculty of Health Sciences, Universitat Jaume I, Castellón de la Plana, Castellón, Spain

Francisco Javier Soriano-Vidal^{a,b,c,d}

^aDepartment of Obstetrics and Gynaecology, Hospital Luis Alcanyis, Avda. Ausiàs March, 46800 Xàtiva, Spain

^bFoundation for the Promotion of Health and Biomedical Research in the Valencian Region (FISABIO), Valencia, Spain

^cDepartment of Nursing, Universitat de València, C/Jaume Roig, s/n, 46001 Valencia, Spain

^dDepartment of Nursing, University of Alicante, Cta. San Vicente del Raspeig s/n 03690 San Vicente del Raspeig, Alicante, Spain

Rafael Vila-Candel^{a,b,c}

^aDepartment of Obstetrics and Gynaecology, Hospital Universitario de la Ribera, Crta. Corbera km 1, 46.600 Valencia, Spain

^bFoundation for the Promotion of Health and Biomedical Research in the Valencian Region (FISABIO), Valencia, Spain

^cDepartment of Nursing, Universitat de València, C/Jaume Roig, s/n, 46001 Valencia, Spain

* Corresponding author.

E-mail addresses: e.castro-sanchez@imperial.ac.uk (E. Castro-Sánchez),
dmena@uji.es (D. Mena-Tudela),
soriano.fravid@gva.es (F. Soriano-Vidal),
rafael.vila@ucv.es (R. Vila-Candel).

Received 24 April 2020