

Sewage-based Epidemiology Requires a Truly Transdisciplinary Approach

If asked whether you had consumed illicit drugs recently, would you admit it? If yes, could you precisely recall types of drug, times and amounts used? If you were the person commissioned with the task of quantifying drug use, what approach

would you use given the social stigma attached with such behavior? We measure drug residues in sewage, which represents urine of entire populations, to provide an objective estimate of total drug use in a region.

In transdisciplinary projects, sewage-based results provide valuable information at unrivaled spatiotemporal resolution complementing traditional data.

Christoph Ort, Caleb J. Banta-Green, Lubertus Bijlsma, Sara Castiglioni, Erik Emke, Coral Gartner, Barbara Kasprzyk-Hordern, Malcolm J. Reid, Jörg Rieckermann, Alexander L. N. van Nuijs

Sewage-based Epidemiology Requires a Truly Transdisciplinary Approach | GAIA 23/3 (2014): 266–268

Keywords: drugs of abuse, illicit drugs, public health, sewage, transdisciplinarity

Context and Transdisciplinarity

Directly and indirectly, illicit drug use causes substantial global health problems and social harms (e. g., overdose, blood borne viruses, violence) (Degenhardt and Hall 2012). Accurate and reliable data on illicit drug consumption is needed to plan and assess the impact of interventions aimed at reducing drug use. Traditional monitoring methods that rely on self-report (e. g., population surveys) may underestimate drug use because it is an illegal, stigmatized activity. Additionally, these methods are expensive, time-consuming and incomplete (Banta-Green and Field 2011). Therefore, scientists have developed an innovative, complementary method of estimating community drug use by quantifying drug residues in sewage, which contains urine from the entire population: they 1. collect a sewage sample, 2. measure concentrations of drug residues in the samples, 3. calculate sewage drug loads (concentrations multiplied by sewage volume), 4. correct for human metabolism (excretion rate), and 5. normalize total loads with population size.

To ensure reliable results, sewage-based epidemiology (SBE) requires significant expertise from numerous disciplines: environmental engineers design sample collection to guarantee that

sewage samples are representative of the community's excreted urine; analytical chemists develop methods to accurately quantify drug residues; environmental chemists investigate transformation in sewers; pharmacologists define the identity and amounts of excreted drug residues to estimate total consumption; epidemiologists identify how this information can complement traditional methods to estimate prevalence of substance use; and social scientists assess ethical aspects of sewage analysis.

This paper does not describe the outcomes of a specific project, but emphasizes the beneficial collaboration of individuals with a mutual interest, at various levels – without the involvement of specific funding and virtually no extra administration.

Sewage Systems, Sampling and Chemical Analysis

Flushed via toilets, excreted drug residues in sewers can be subject to high short-term dynamics and diurnal variations. Both need to be considered to obtain a representative sample (Ort et al. 2010). A thorough experimental design assesses the definition of catchment area, hydraulic properties and the sampling setup, which are elicited from utilities with specifically tailored questionnaires (Castiglioni et al. 2013). Typically, a sample of 500 milliliters of raw sewage, filtered and stabilized until analysis, is sufficient. Analyte concentrations are at the low nanogram per liter level which necessitates sample clean-up, pre-concentration and subsequent chemical analysis via highly specific and sensitive techniques, normally based on liquid chromatography coupled to tandem mass spectrometry (Van Nuijs et al. 2011).

Contact: Dr. Christoph Ort | Swiss Federal Institute of Aquatic Science and Technology (Eawag) | Urban Water Management | Ueberlandstr. 133 | 8600 Dübendorf | Switzerland | Tel.: +41 58 7655277 | E-Mail: christoph.ort@eawag.ch

© 2014 C. Ort et al.; licensee oekom verlag. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Back-calculation and Uncertainty

The analyte of choice in SBE is generally the primary urinary metabolite of the drug of interest. To differentiate between actual consumption and direct disposal of unused drugs, enantiomeric profiling can be applied with the usage of chiral chromatography coupled to mass spectrometry (Kasprzyk-Hordern and Baker 2012). To facilitate comparison among cities and to estimate consumption, sewage drug loads are normalized by the number of people and corrected for excretion rates of the metabolite and drug purity. Excretion rates are available from clinical studies, and drug purity is determined through analysis of drugs from local seizures or test purchases. Interestingly, a reliable estimate of the population contributing to the sewage is difficult: census may be outdated or not coincide with the geographical boundary of a sewer catchment, and accounting for commuters and tourists is challenging. One solution is to measure simultaneously specific compounds in the sewage samples to estimate population size. Ideally, a ubiquitous biomarker could be found that has no other source than human excretion and is stable during transport in sewers (Chen et al. 2014). So far, pharmaceutical residues have been successfully combined with Bayesian inference (O'Brien et al. 2014). Although transformation of certain drugs during transport in sewers may affect SBE, this is just starting to be investigated (Thai et al. 2014).

In most cases, sewage-based estimates of substance use – as pure substance per 1,000 people per day – have an uncertainty of about 40 percent. While this uncertainty may seem large, it is considered acceptable by epidemiologists as there is lower risk of the estimates being biased. By contrast, other monitoring methods (e.g., surveys) may have narrow variability components, but higher risk of bias. Currently, the estimation of the number of doses or even of individual users based on sewage samples requires a series of assumptions and remains speculative. Nevertheless, some studies have shown good qualitative agreement between sewage-based estimates and other data (Reid et al. 2012).

Applications, Ethical Considerations and Outlook

Most studies aimed at regional or national estimates of drug use, but international comparisons exist now (see figure). Sewage analysis can be carried out at high temporal frequency and spatial resolution. Daily variations can be identified (e.g., a weekend-effect for cocaine and ecstasy), and intra-day variability might discriminate between commuters and permanent residents (Brewer et al. 2012). The approach also allows investigating holiday periods, special events in sports or music with permission from the organizers (Lai et al. 2013). Given the negligible risk of (in)direct harm




FIGURE: Population-normalized ecstasy (MDMA) sewer loads during two one-week periods (seven daily samples each) in 28 sewage treatment plant catchments (adapted from Ort et al. 2014).

to residents, minimal ethical issues are expected for SBE in large, general populations ($\geq 10,000$ people) if findings do not relate to specific groups (e. g., a disadvantaged suburb) (Hall et al. 2012).

SBE is not limited to illicit drugs. Examples encompass anti-histamine use in the general population (Reid et al. 2011) or Attention Deficit Hyperactivity Disorder medications used by students during semester and examination periods (Burgard et al. 2013). Identifying doping used by athletes was suggested but may prove difficult due to sampling issues and other practical aspects (Harman et al. 2011). It is expected that SBE can expand to further public health aspects (e. g., lifestyle and diet) (Thomas and Reid 2011), and may serve as an early warning system in case of pandemics.

Collaboration with various stakeholders from, for example, public health and drug prevention institutes, forensics and law enforcement is now envisaged to apply SBE meaningfully in a local, relevant context. The innovative co-production of knowledge requires diverse perspectives from an early project stage onwards to reach the desired societal impact.

We would like to express our gratitude to colleagues who advanced SBE. They cover a wide range of disciplines, from psychologists to chemists and statisticians to toxicologists, medical doctors and many more. Thank you Will Backe, Frédéric Béen, Jean-Daniel Berset, Alex Brewer, Raimondo Bruno, Steve Carter, Adrian Covaci, Pim de Voogt, Olivier Delémont, Geoff Eaglesham, Jonas Maria Eppler, Pierre Esseiva, Jennifer Field, Wayne Hall, Christopher Harman, Félix Hernández, Martina Kinzig, Paul Kirkbride, Foon Yin Lai, Michael Lawrence, Jochen Mueller, Jake O'Brien, Jeremy Prichard, Andreas Scheidegger, Fritz Sörgel, Phong Thai, Kevin Thomas, Peter Vallely and Ettore Zuccato.

References

- Banta-Green, C. J., J. A. Field. 2011. City-wide drug testing using municipal wastewater: A new tool for drug epidemiology. *Significance* 8/2: 70–74.
- Brewer, A. J., C. Ort, C. J. Banta-Green, J.-D. Berset, J. A. Field. 2012. Normalized diurnal and between-day trends in illicit and legal drug loads that account for changes in population. *Environmental Science and Technology* 46/15: 8305–8314.
- Burgard, D. A., R. Fuller, B. Becker, R. Ferrell, M. J. Dinglasan-Panlilio. 2013. Potential trends in Attention Deficit Hyperactivity Disorder (ADHD) drug use on a college campus: Wastewater analysis of amphetamine and ritalinic acid. *Science of the Total Environment* 450–451: 242–249.
- Castiglioni, S. et al. 2013. Evaluation of uncertainties associated with the determination of community drug use through the measurement of sewage drug biomarkers. *Environmental Science and Technology* 47/3: 1452–1460.
- Chen, C., C. Kostakis, J. P. Gerber, B. J. Tscharke, R. J. Irvine, J. M. White. 2014. Towards finding a population biomarker for wastewater epidemiology studies. *Science of the Total Environment* 487: 621–628.
- Degenhardt, L., W. Hall. 2012. Extent of illicit drug use and dependence, and their contribution to the global burden of disease. *Lancet* 379/9810: 55–70.
- Hall, W. et al. 2012. An analysis of ethical issues in using wastewater analysis to monitor illicit drug use. *Addiction* 107/10: 1767–1773.
- Harman, C., M. Reid, K. V. Thomas. 2011. Concerning the viewpoint; “an anti-doping sampling strategy utilizing the sewerage systems of sport villages”. *Environmental Science and Technology* 45/10: 4191.
- Kasprzyk-Hordern, B., D. R. Baker. 2012. Estimation of community-wide drugs use via stereoselective profiling of sewage. *Science of the Total Environment* 423: 142–150.
- Lai, F. Y. et al. 2013. Using quantitative wastewater analysis to measure daily usage of conventional and emerging illicit drugs at an annual music festival. *Drug and Alcohol Review* 32/6: 594–602.
- O'Brien, J. W. et al. 2014. A model to estimate the population contributing to the wastewater using samples collected on census day. *Environmental Science and Technology* 48/1: 517–525.
- Ort, C., M. G. Lawrence, J. Rieckermann, A. Joss. 2010. Sampling for Pharmaceuticals and Personal Care Products (PPCPs) and illicit drugs in wastewater systems: Are your conclusions valid? A critical review. *Environmental Science and Technology* 44/16: 6024–6035.
- Ort, C. et al. 2014. Spatial differences and temporal changes in illicit drug use in Europe quantified by wastewater analysis. *Addiction* 109/8: 1338–1352.
- Reid, M. J., K. H. Langford, J. Mørland, K. V. Thomas. 2011. Quantitative assessment of time dependent drug-use trends by the analysis of drugs and related metabolites in raw sewage. *Drug and Alcohol Dependence* 119/3: 179–186.
- Reid, M. J. et al. 2012. Estimation of cocaine consumption in the community: A critical comparison of the results from three complimentary techniques. *BMJ Open* 2/6: e001637. doi:10.1136/bmjopen-2012-001637.
- Thai, P. K., G. Jiang, W. Gernjak, Z. Yuan, F. Y. Lai, J. F. Mueller. 2014. Effects of sewer conditions on the degradation of selected illicit drug residues in wastewater. *Water Research* 48/1: 538–547.
- Thomas, K. V., M. J. Reid. 2011. What else can the analysis of sewage for urinary biomarkers reveal about communities? *Environmental Science and Technology* 45/18: 7611–7612.
- Van Nuijs, A. L. N. et al. 2011. Illicit drug consumption estimations derived from wastewater analysis: A critical review. *Science of the Total Environment* 409: 3564–3577.

Submitted May 26, 2014; revised version accepted July 9, 2014.

CONTRIBUTING AUTHORS

Dr. Christoph Ort

Dr. Jörg Rieckermann

both: Swiss Federal Institute of Aquatic Science and Technology (Eawag), Dübendorf, Switzerland

Dr. Caleb J. Banta-Green

University of Washington, Alcohol and Drug Abuse Institute, Seattle, WA, United States

Lubertus Bijlsma, MSc

University Jaume I, Research Institute for Pesticides and Water, Castellón de la Plana, Spain

Dr. Sara Castiglioni

IRCCS – Istituto di Ricerche Farmacologiche Mario Negri, Department of Environmental Health Sciences, Milan, Italy

Erik Emke, BSc

KWR Watercycle Research Institute, Nieuwegein, The Netherlands

Dr. Coral Gartner

The University of Queensland, UQ Centre for Clinical Research, Royal Brisbane and Women's Hospital, Herston, QLD, Australia

Dr. Barbara Kasprzyk-Hordern

University of Bath, Department of Chemistry, Bath, United Kingdom

Malcolm J. Reid, BSc (Hons)

Norwegian Institute for Water Research (NIVA), Oslo, Norway

Dr. Alexander L. N. van Nuijs

University of Antwerp, Toxicological Center, Antwerp, Belgium

Christoph Ort



Born 1974 in Zurich, Switzerland. 2006 PhD in environmental engineering (ETH Zurich). 2008 to 2010 postdoc at the Advanced Water Management Centre (University of Queensland, Australia). Since 2011 group leader in the Urban Water Management Department (Eawag), Dübendorf, Switzerland. Research interests: water-related urban pollutant fluxes, modeling, sewer processes, sampling and monitoring design, source typing and tracking.