

Dramatic Effect of the Gas Atmosphere on the Deprotection of (Z)- γ - Hydroxy- α,β -unsaturated Esters

Irakusne López[†], Santiago Rodríguez^{*†}, Javier Izquierdo[†], Florenci V. González^{*†} and Cristian Vicent[‡]

*Departament de Química Inorgànica i Orgànica and Serveis Centrals d'Instrumentació Científica, Universitat
Jaume I, 12080 Castelló, Spain*

Abstract-The deprotection of *O*-protected (*Z*)- α -methyl- γ -hydroxy- α,β -unsaturated ester **1** furnishes dimer **2** when the reaction is performed under a nitrogen atmosphere, while compound **3** forms when the reaction is performed under an oxygen atmosphere.


Keywords: butenolides, dimerization, γ -hydroxybutenolides

1. Introduction

Butenolides are interesting building blocks in organic synthesis [1], and their dimerization offers an attractive approach for the synthesis of natural products [2]. The formation of $\alpha,\beta,\gamma,\delta$ dimers from butenolides under basic conditions has been reported [3,1b], and the γ -hydroxybutenolide derivatives are known antimicrobial compounds and are often used as anticancer or anti-inflammatory agents [4]. Thus, the development of facile methods for synthesizing this class of structures is important [5,2d]. Herein, we report that both types of compounds ($\alpha,\beta,\gamma,\delta$ dimers and γ -hydroxybutenolides) can be obtained from the same butenolide, depending upon the reaction conditions. Under an inert atmosphere, the $\alpha,\beta,\gamma,\delta$ dimer is formed, but under an oxygen atmosphere the γ -hydroxybutenolide is obtained.


2. Results

In the course of our work on the epoxidation of γ -hydroxy- $\alpha,\beta,\gamma,\delta$ -unsaturated esters [6], we easily prepared *O*-protected (*Z*)- α -methyl- γ -hydroxy- $\alpha,\beta,\gamma,\delta$ -unsaturated ester **1** from (*S*)-lactaldehyde [7]. During the deprotection of (*Z*) enoate **1**, we observed the formation of dimer **2**. Serendipitously, when the deprotection reaction was carried out in the open air, compound **3** was obtained. Presumably, deprotection of compound **1** gave rise to γ -hydroxy enoate **4**, which, upon cyclization, furnished butenolide **5**. Butenolide **5** would subsequently be converted into dimer **2** or, if the oxidative process takes place promoted by an oxygen atmosphere, compound **3** (Scheme 1).




Scheme 1.

The formation of dimer **2** can be easily explained by the basic enolization of butenolide **5**, giving rise to enolate **6** which, upon conjugate addition with **5**, furnishes dimer **2** (Scheme 2). Compound **2** was formed in a highly stereoselective fashion (d.r. 19:1) giving rise to the *endo*-product. It is noteworthy that 4 stereocenters were formed in one pot with stereoselectivity. Although Michael dimerizations of α -substituted butenolides [3] have been previously reported, we report here a stereoselective Michael dimerization of a $\alpha\beta\gamma\delta$ -disubstituted butenolide. Thus, Michael dimerization is more selective toward $\alpha\beta\gamma$ -disubstituted butenolides than toward α -substituted butenolides.



Scheme 2.

Stereochemical assignment in compound **2** was performed by NOE experiments. Dimer **2** gave NOE between H-4' and H-2' and between H-4' and H-4, while H-2' also gave NOE with H-4 (Scheme 3).



Scheme 3.

The X-ray crystal structure of **2** definitively confirmed the stereochemistry (Figure 1). Compound **2** crystallizes in the centrosymmetric $C2/c$ space group. Figure 1a shows the molecular structure of the corresponding enantiomers related by a crystallographic inversion center. The structure of **2** comprises two linked five-membered lactones. Both rings are almost coplanar (interplanar angle below 10°) and their relative orientation is linear with respect to the long axis of the 5-membered ring crossing at the C linking both lactones.





Fig. (1).

In contrast, compound **3** was obtained when the deprotection reaction was performed under an oxygen atmosphere. The α -substituted α -hydroxybutenolides are known to form through photooxidation of furans [8]. However, formation of compound **3** takes place without any photosensitizer, so singlet oxygen is not generated. In a

mechanism similar to the autoxidation of benzylic carbanions [9], carbanion **6** combines with molecular oxygen, furnishing intermediate **7** which, upon reaction with another molecule of starting enolate **6**, gives hemiketal **3**. The molecular structure of compound **3** has also been unambiguously determined by single-crystal X-ray methods (see Figure 1 b).



Scheme 4.

3. Conclusions

In summary, we have shown that the deprotection of enoate **1** furnishes □□□' dimer **2** in a highly stereoselective fashion if the reaction is performed under a nitrogen atmosphere whilst deprotection of enoate **1** under an oxygen atmosphere furnishes □-hydroxybutenolide **3**.

4. Experimental Section

General Experimental Methods. All solvents used in reactions were freshly distilled from appropriate drying agents before use. ^1H NMR spectra and ^{13}C NMR spectra were measured in CDCl_3 (^1H , 7.24 ppm; ^{13}C 77.0 ppm) solution at 30 °C on a 300 MHz Mercury Varian or a 500 MHz Innova Varian NMR spectrometer at the Serveis Centrals d'Instrumentació Científica de la Universitat Jaume I. Mass spectra were measured in a QTOF I (quadrupole-hexapole-TOF) mass spectrometer with an orthogonal Z-spray-electrospray interface (Micromass, Manchester, UK). IR spectra were recorded as oily films on NaCl plates on a Perkin-Elmer 2000 FT-IR spectrometer. EM Science Silica Gel 60 was used for column chromatography while TLC was performed with E. Merck precoated plates (Kieselgel 60, F₂₅₄, 0.25 mm). Unless otherwise specified, all reactions were carried out under argon atmosphere with magnetic stirring. About the X-ray crystallographic study, the crystals are air stable and were mounted on the tip of a glass fiber with the use of epoxi cement. X-ray diffraction experiment was carried out on a Bruker SMART CCD diffractometer using Mo-K α radiation ($\lambda = 0.71073 \text{ \AA}$) at room temperature. The data were collected with a frame width of 0.3 ° in ω and a counting time of 30 and 60 s per frame for compounds **2** and **3** at a crystal to detector distance of 4 cm. The diffraction frames were integrated using the SAINT package and corrected for absorption with SADABS [10]. The structures were solved by direct methods and refined by the full-matrix method based on F^2 using the SHELXTL software package [11]. All non hydrogen atoms were refined anisotropically. Hydrogen atoms were generated geometrically, assigned isotropic thermal parameters and allowed to ride on their respective parent carbon atoms.

(Z)-Ethyl 4-(tert-Butyl-dimethyl-silanyloxy)-2-methyl-pent-2-enoate 1. To an ice-bath cold suspension of sodium hydride (60% in mineral oil) (2.39 g, 59.75 mmol) in THF (59 mL) was added triethyl 2-phosphonopropionate (13 mL, 59.42 mmol). The resulting mixture was stirred at room temperature for 1h and then was cooled with an ice-bath and a solution of *O*-tert-butyldimethylsilyl lactaldehyde [12] (10.21 g, 54.30 mmol) in THF (40 mL) was added. The resulting mixture was stirred at room temperature for 7h and then was quenched with brine and extracted with Et_2O (3 x 30 mL), the organic layers were washed (brine), dried (Na_2SO_4) and concentrated. The crude oil was purified by silica gel chromatography, eluted with hexanes/ Et_2O (99:1) to afford 5.85 g of compound **1** and 6.8g of *E* isomer (86%). $[\alpha]_{\text{D}}^{20} = -5.3$ ($c = 1.3$, CHCl_3).

Spectroscopic data for **1**: ^1H NMR (500MHz, CDCl_3) δ 5.91 (1H, dd, $J=8.0$ and 1.5Hz), 5.15 (1H, dq, $J=7.0\text{Hz}$), 4.19 (2H, m), 1.88 (3H, d, $J=1.5\text{Hz}$), 1.30 (3H, t, $J=7.0\text{Hz}$), 1.22 (3H, d, $J=6.5\text{Hz}$), 0.87 (9H, s), -0.05 (3H, s), -0.06 (3H, s). ^{13}C NMR (125 MHz, CDCl_3) δ 167.8, 148.1, 125.1, 66.7, 60.7, 26.2, 24.1, 20.5, 18.5, 14.6, -4.3, -4.4 ppm. IR (NaCl) δ 2957, 2930, 2858, 1719, 1591, 1473, 1374, 1255, 1126, 1078, 1005, 835, 776 cm^{-1} . HRMS m/z calcd. for $\text{C}_{14}\text{H}_{28}\text{O}_3\text{SiNa}$ [$\text{M}+\text{Na}^+$]: 295.1705, found: 295.1689.

(R)-5-((2R,3S,4S)-tetrahydro-2,4-dimethyl-5-oxofuran-3-yl)-3,5-dimethylfuran-2(5H)-one 2. To an ice cold solution of compound **1** (2.65 g, 9.74 mmol) in THF (28 mL) under a nitrogen atmosphere was added tetra-*n*-butylammonium fluoride (1.0 M in THF) (19 mL, 19 mmol). The resulting mixture was stirred under nitrogen and allowed to warm to room temperature over 16 h. The reaction then was quenched with brine and extracted with Et_2O (3 x 30 mL), and the organic layers were washed (brine), dried (Na_2SO_4) and concentrated. The crude oil was purified by silica gel chromatography, eluted with hexanes/ EtOAc (1:1) and ethyl acetate, to afford 1.63 g (75%) of compound **2** as a white solid (m.p. 68-69°C).

Spectroscopic data for **2**: ^1H NMR (500 MHz, CDCl_3) δ 6.99 (1H, s), 4.22 (1H, dd, $J=6.6$, 6.3Hz), 2.64 (1H, dd, $J=7.5$, 8.4Hz), 2.10 (1H, dd, $J=7.2$, 8.4Hz), 1.97 (3H, d, $J=1.8\text{Hz}$), 1.52 (3H, s), 1.44 (3H, d, $J=5.7\text{Hz}$), 1.42 (3H, d, $J=6.9\text{Hz}$). ^{13}C NMR (125 MHz, CDCl_3) δ 177.6, 172.3, 150.2, 131.0, 85.1, 74.7, 57.1, 37.5, 23.0, 22.7, 18.1, 10.6 ppm. IR (NaCl) δ 2982, 1770, 1456, 1194, 893, 765, 713 cm^{-1} . HRMS m/z calcd. for $\text{C}_{12}\text{H}_{16}\text{O}_4\text{Na}$ [$\text{M}+\text{Na}^+$]: 247.0946, found: 247.0958.

5-hydroxy-3,5-dimethylfuran-2(5H)-one 3. An oxygen-flushed mixture of compound **1** (1.5 g, 5.51 mmol) in THF (28 mL) was cooled with an ice bath. Then tetra-*n*-butylammonium fluoride (1.0 M in THF) (11 mL, 11 mmol) was added dropwise. The resulting mixture was stirred under oxygen atmosphere at room temperature for 3 h. The reaction was quenched with brine, the organic solvent was removed under vacuum and then extracted with Et_2O (3 x 30 mL). The organic layers were washed (brine), dried (Na_2SO_4), and concentrated. The crude oil was purified through chromatography (silica-gel, hexanes/ EtOAc (6:4), (1:1) and ethyl acetate) to afford 494 mg (70%) of compound **3** as a white solid (m.p. 98-99°C) (lit. 98-99°C) [13].

Spectroscopic data for **3**: ^1H NMR (500 MHz, CDCl_3) δ 6.85 (2H, s), 1.88 (6H, d, $J=1.8\text{Hz}$), 1.66 (6H, s). ^{13}C NMR (125MHz, CDCl_3) δ 172.3, 148.1, 131.5, 104.6, 24.7, 10.3 ppm. IR (NaCl) δ 3278, 2926, 1741, 1450, 1192, 1049, 929, 873, 713 cm^{-1} . HRMS m/z calcd. for $\text{C}_6\text{H}_8\text{O}_3\text{Na}$ [$\text{M}+\text{Na}^+$]: 151.0371, found: 151.0365.

5. Acknowledgments

This work was financed by Bancaixa-UJI foundation (P1 1A2005-14).

6. References and Footnotes

* Corresponding autor: Tel. 34 964729156; fax. 34 964728214 ; e-mail: fgonzale@qio.uji.es

† Departament de Química Inorgànica i Orgànica, Universitat Jaume I

‡ Serveis Centrals d'Instrumentació Científica, Universitat Jaume I

[1] a) Rao, Y.S. *Chem. Rev.* **1964**, *64*, 353-388. b) Rao, Y.S. *Chem. Rev.* **1976**, *76*, 625-694.

[2] a) Bagal, S. K.; Adlington, R. B.; Baldwin, J. E.; Marquez, R.; Cowley, A. *Org. Lett.* **2003**, *17*, 3049-3052. b) Bagal, S. K.; Adlington, R. B.; Marquez, R.; Cowley, A.; Baldwin, J. E. *Tetrahedron Lett.* **2003**, *44*, 4993-4996. c) Bagal, S. K.; Adlington, R. B.; Baldwin, J. E.; Marquez, R. *J. Org. Chem.* **2004**, *69*, 9100-9108. d) Bagal, S. K.; Adlington, R. B.; Brown, R. A. B.; Baldwin, J. E. *Tetrahedron Lett.* **2005**, *46*, 4633-4637.

[3] Okano, T.; Chokai, M.; Eguchi, S.; Hayakawa, Y. *Tetrahedron* **2000**, *56*, 6219-6222.

[4] a) Wright, A. D.; de Nys, R.; Angerhofer, C. K.; Pezzuto, J. M.; Gurrath, M. *J. Nat. Prod.* **2006**, *69*, 1180-1187. b) Keyzers, R. A.; Davies-Coleman, M. T. *Chem. Soc. Rev.* **2005**, *34*, 355-365. c) Grossman, G.; Poncioni, M.; Bornand, M.; Jolivet, B.; Neuburger, M.; Sequin, U. *Tetrahedron* **2003**, *59*, 3237-3251. d) Charan, R. D.; McKee, T. C.; Boyd, M. R. *J. Nat. Prod.* **2001**, *64*, 661-663.

[5] a) Brückner, R. *Curr. Org. Chem.* **2001**, *5*, 679-718. b) Carter, N. B.; Nadany, A. E.; Sweeney, J. B. *J. Chem. Soc., Perkin Trans.* **2002**, *1*, 2324-2342. c) Patil S. N.; Liu, F. *J. Org. Chem.* **2008**, *73*, 4476-4483.

[6] López, I.; Rodríguez, S.; Izquierdo, J.; González, F.V. *J. Org. Chem.* **2007**, *62*, 6614-6617.

- [7] Compound **1** is prepared from *O*-protected (*S*)-lactaldehyde by a Horner-Emmons reaction using triethyl 2-phosphonopropionate, furnishing a 3:2 mixture of *E/Z* isomers easily separable through chromatography (see ref. 6 and experimental section).
- [8] a) Patil S. N.; Liu, F. *Org. Lett.* **2007**, *9*, 195-198. b) Patil S. N.; Liu, F. *J. Org. Chem.* **2007**, *72*, 6305-6308. and ref. 5c and cites herein.
- [9] a) Russell, G. A.; Janzen, E. G.; Becker, H-D.; Smentowski, F. J. *J. Am. Chem. Soc.* **1962**, *84*, 2652-2653. b) Russell, G. A.; Bemis, A. G. *J. Am. Chem. Soc.* **1966**, *88*, 5491-5497. c) Barton, D. H. R.; Jones, D. W. J. *J. Chem. Soc.* **1965**, 3563-3570.
- [10] a) *SAINT*; 5.0 ed.; Bruker Analytical X-Ray Systems: Madison, WI, **1996**. b) Sheldrick, G. M., *SADABS empirical absorption program*; University of Göttingen, **1996**.
- [11] Sheldrick, G. M., *SHELXTL*; 5.1 ed.; Bruker Analytical X-Ray Systems: Madison, WI., **1997**.
- [12] The preparation of *O*-*tert*-butyldimethylsilyl lactaldehyde was performed according to: Marshall, J.A.; Shiping, X. *J. Org. Chem.* **1995**, *60*, 7230-7237.
- [13] Haval, Kishan P.; Argade, Narshinha P. *Synthesis* **2007**, *14*, 2198 - 2202.