

An Adaptive Virtual Reality System for the Treatment of Adjustment Disorder and Complicated Grief: 1-year Follow-up Efficacy data

Soledad Quero^{a,c*}, Mar Molés^a, Daniel Campos^a, Sabrina Andreu-Mateu^a, Rosa M. Baños^{b,c},
Cristina Botella^{a,c}

^aDepartamento de Psicología Básica, Clínica y Psicobiología, Universitat Jaume I; Castellón, Spain

^bDepartamento de Personalidad, Evaluación y Tratamientos Psicológicos, Universidad de Valencia, Valencia,
Spain

^cCIBER de Fisiopatología de la Obesidad y Nutrición (CIBERON), Instituto Salud Carlos III, Madrid, Spain

Abstract: Adjustment disorders (AD) and complicated grief (CG) are serious mental conditions that have a high prevalence and are associated with significant impairments in social and work functioning. Recently, these categories have been better specified in the new ICD-11 proposal. Empirical research on the efficacy of treatments for these problems is scarce. This study aims to offer long-term efficacy data from a between-groups controlled study that compares two treatment conditions (AD-protocol applied in a traditional way: N= 18; and the same protocol supported by virtual reality-VR-: N =18) and a waiting list (WL) control group (N =18). Both treatment conditions resulted in statistically significant improvements on both primary and secondary outcome measures, with large effect sizes, and this improvement did not occur in the WL. These changes were maintained in both treatment conditions in the medium (6-month) and long term (12-month follow-up). Larger effect sizes were achieved in the VR condition in the long term. Furthermore, clinically significant change estimations on the primary outcome measures showed an advantage for the VR condition. This is the first controlled study to compare a traditional face-to-face protocol for the treatment of stress-related disorders to the same protocol supported by VR and a WL control group.

Key Practitioner Message:

- A CBT protocol has been designed and developed for AD.
- This protocol applied traditionally and with VR was more effective than a WL.
- Changes were maintained in both treatment conditions at 6- and 12-month follow-ups.
- VR seems to be a promising tool to deliver the elaboration/exposure component for AD and CG.

Key words: Adjustment Disorder, Complicated Grief, Prolonged Grief Disorder, CBT, Virtual reality, Positive psychology strategies.

* Corresponding author:

Soledad Quero PhD.

Departamento de Psicología Básica, Clínica y Psicobiología. Universitat Jaume I.

Avda. Vicente Sos Baynat s/n 12071 Castellón (Spain)

Phone: +34 964387641

Fax: +34 964729267

E-mail: squero@uji.es

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/cpp.2342

1. INTRODUCTION

Adjustment disorders (AD) and complicated grief (CG) are two categories that have traditionally been poorly specified by the DSM and ICD-10. In the last edition (DSM-5, APA, 2013), AD was classified for the first time in the category of “Trauma and stress related disorders”, along with another important disorder, posttraumatic stress disorder (PTSD). However, the DSM-5 has been criticized for the lack of specific symptom descriptions and the difficulty of distinguishing between AD and normal adaptive stress reactions (Casey, 2014). In the case of CG, DSM-5 has introduced criteria for persistent complex bereavement disorder also categorized as one of the “other specified trauma- and stressor-related disorder” while the CG, was placed in the Research section (III) as a condition requiring further study.

More recently, the ICD-11 work group proposed that these two diagnostic categories to be included for the first time in the new chapter “Disorders Specifically Associated with Stress” and have provided a clearer definition of symptoms (Maercker et al., 2013). AD is defined as a maladaptive reaction to a negative stressful life event or life change, characterized by symptoms of preoccupation, such as excessive worry, recurrent and distressing thoughts about the stressor, or constant rumination about its implications. There is a failure to adapt, i.e., the symptoms interfere with everyday functioning, such as difficulties concentrating or sleep disturbances resulting in performance problems. These symptoms can also be associated with loss of interest in work, social life, caring for others, leisure activities, resulting in impairments in social or occupational functioning. Prolonged grief disorder is defined as a disturbance in which, following the death of a person close to the bereaved, there is a persistent and pervasive yearning or longing for the deceased, or a persistent preoccupation with the deceased that extends for an abnormally long period beyond the expected social and cultural norms (e.g., at least 6 months, or longer, depending on cultural and contextual factors) and is sufficiently severe to cause significant impairment in the person’s functioning. The response can also be characterized by difficulties in accepting the death, feeling that one has lost a part of one’s self, anger about the loss, guilt, or difficulty in engaging with social or other activities. In this article, we use the term CG to refer to the Prolonged Grief disorder proposed by the ICD-11. Participants with persistent complex bereavement disorder were not included (Boelen & Smid, 2017).

However, at the time the present study was conducted, the classifications mentioned above were not available. Therefore, the DSM-IV-TR (APA, 2000) and ICD-10 (WHO, 1992) criteria for AD were followed. In the case of CG, according to the ICD-10 classification, we considered it as an AD. Specifically, the ICD-10 allows the codification of

bereavement as an AD when the reaction to the death of a loved one is considered abnormal because of its manifestations or content, or when this reaction is still intense more than 6 months after the death.

AD is among the diagnoses most widely used by psychiatrists and psychologists worldwide (Maercker *et al.*, 2013). Recently, Yaseen (2017) estimated AD prevalence data in outpatient psychiatric clinics in values around 11%. Regarding CG, a recent systematic review and meta-analysis on the prevalence of prolonged grief disorder in adult bereavement conducted by Lundorff, Holmgren, Farver-Vestergaard and O'Connor. (2017) revealed a pooled prevalence of 9.8%. In addition to the high prevalence of AD (Carta, Balestrieri, Murru & Hardoy, 2009), this problem is associated with significant impairment in social and work functioning, causing a high percentage of sick leave (Van der Klink, Blonk, Schene & van Dijk, 2003). Furthermore, AD is the most common diagnosis in people with self-harm behaviors (Casey, 2009); up to 25% of adolescents with AD perform suicidal behaviors (Pelkonen, Marttunen, Henriksson & Loongvist, 2007), with this percentage reaching 60% in adults (Kryzhananovskaya & Canterbury, 2001). Results obtained more recently by Casey, Jabbar, O'Leary and Doherty (2015) indicate that AD is a potentially serious condition that can present life-threatening features, with similar percentages of patients with AD and depressive episodes reporting suicidal ideation or behavior. In addition, the occurrence of suicidality at a lower symptom score in AD suggests that this group is more vulnerable. Along the same lines, CG is associated with intense suffering and the risk of developing mental and physical health problems (Stroebe, Schut & Stroebe, 2007), increasing the risk of hospitalization (Li, Laursen, Precht, Olsen, & Mortensen, 2005), and being associated with medical comorbidity (Prigerson *et al.*, 1997), suicidality (Latham & Prigerson, 2004), and greater work and social impairment. It is a debilitating and chronic disorder (Simon *et al.*, 2007).

Therefore, evidence-based psychological treatments for AD and CG are needed. The few studies conducted using between-group designs for AD have been carried out from different therapeutic approaches: mirror therapy (González-Jaimes & Turnbull-Plaza, 2003), problem solving and temporal contingency approach (van der Klink *et al.*, 2003), CBT (Sterinhardt & Dolbier; 2008), brief group psychodynamic therapy (Ben-Itzhak *et al.*, 2012), Body-Mind-Spirit psychotherapy (Hsiao *et al.*, 2014), meditation training (Srivastava, Talukdar & Lahan, 2011). Although overall they reported a decrease of symptoms, many of them have important methodological limitations. Regarding treatment for CG, Neimeyer has a number of publications about treatments for this problem. His approach (Neimeyer, 2000)

focuses on the reconstruction of meaning after loss (Neimeyer & Currier, 2009). Results from the 11 studies included in the review study by Currier, Holland and Neimeyer (2010) showed that CBT-based interventions were more effective than other commonly practiced therapies. However, after accounting for the influence of researcher allegiance, differences between therapies became smaller and were no longer statistically significant. Compared to no-treatment control groups, CBT-based interventions produced benefits immediately after intervention, but they did not yield statistically significant overall effects at follow-up. Therefore, the authors concluded that there is preliminary evidence for the helpfulness of CBT-based interventions for bereaved persons, but studies on the relative efficacy of different cognitive-behavioral change strategies, as well as other orientations, are still needed. As indicated in more recent review studies (Shear, 2015; Crunk, Kurke & Robinson III, 2017), several randomized controlled trials have shown that a short-term approach called complicated grief treatment is the treatment that has been most extensively studied so far (e.g., Boelen, Keijsers, van den Hout & van den Hout, 2007; Bryant, Kenny, Joscelyne, et al., 2014; Shear, Frank, Houck & Reynolds, 2005; Shear et al., 2014). The treatment includes two main objectives: restoring effective functioning by generating enthusiasm and creating plans for the future and helping patients find a way to think about the death without having intense feelings of anger, guilt or anxiety (Shear, 2015).

Based on the aforementioned, we can conclude that no specific protocol exists for AD because none of the experimental studies reviewed above propose a treatment specifically designed for this problem. Therefore, we developed an AD-protocol to treat AD and CG (Botella, Baños & Guillén, 2008) that includes CBT techniques and positive psychology strategies. Furthermore, it is supported by virtual reality (VR).

Several systematic reviews (Freeman et al., 2017; Gregg & Tarrier, 2007; McCann et al., 2014; Meyerbroeker & Emmelkamp, 2010) and a number of meta-analytic studies (Opris et al., 2011; Parsons & Rizzo, 2008; Powers & Emmelkamp, 2008) concluded that the most well-established finding is that VR exposure-based treatments can reduce anxiety disorders and PTSD. In the case of Opris et al. (2011) and Morina et al. (2015), the authors also found that VR has a powerful real-life impact, similar to that of the classical evidence-based treatments, and good stability in the results over time. Additionally, meta-analyses and systematic reviews specifically addressed to stress-related disorders (mainly PSD) (Botella, Serrano, Baños & Garcia-Palacios, 2015; Gonçalves, Pedrozo, Coutinho, Figueira & Ventura, 2012; Motraghi, Seim, Meyer & Morissete, 2014), have also shown the usefulness of VR environments for applying the exposure technique in treating these disorders. Finally,

other exposure-based approaches, such as EMDR, have also shown their efficacy in the treatment of distressful experiences that failed to meet the criteria for PTSD (Cvetek (2008).

The present study uses the *EMMA's World* VR system (Botella *et al.*, 2006) to apply elaboration/exposure technique. Preliminary efficacy data for this protocol already exist in two case studies (Andreu-Mateu, Botella, Quero, Guillén & Baños, 2012; Botella, Osma, García-Palacios, Guillén & Baños, 2008), and in a between-group study (Baños *et al.*, 2011) that included participants with different stress-related disorders (PTSD, CG and AD). The results showed that traditional CBT programs were equally as effective as a CBT program supported by *EMMA's World*. However, no control group was included in this study, and no follow-up data were available. The aim of the present study is to offer long-term efficacy data from two AD-treatment protocols (an AD-treatment protocol supported by VR - *EMMA's World*- and an AD-protocol applied in a traditional - face-to face way) compared to a waiting list control group in an RCT. A secondary aim is to explore and compare both ways of delivering the AD-protocol. First, it is hypothesized that both treatment conditions will significantly reduce primary symptoms of AD and CG, compared to the waiting list control group. Second, both treatment conditions will show efficacy and no statistically significant differences will be found between them. Finally, the therapeutic gains obtained in both conditions will be maintained at 1-year follow-up.

2. METHOD

2.1. Design

A three-armed single-blind RCT was conducted. Participants were randomized into three groups: 1) EMMA condition, 2) Traditional condition, and 3) Waiting list (WL) control condition. Participants in the control group were randomly assigned to one of the two treatment conditions after spending time on the waiting list (6 weeks) for ethical reasons. This study was conducted following the CONSORT statement (Consolidated Standards of Reporting Trials, <http://www.consort-statement.org>) and received approval from the Ethics Committee of Universitat Jaume I (Castellón, Spain) Participants were assessed at pre-treatment, post-treatment, and 6- and 12-month follow-ups. Figure 1 shows the flowchart for the study.

2.2. Participants

This study was conducted at the Emotional Disorders Clinic at Universitat Jaume I in Castellón and Valencia (Spain). Participant recruitment was carried out through advertisements (mail, posters, radio, and press) about the study developed in our clinic. Other

participants came to seek help at our clinic, and others were referred from public mental health services.

Inclusion criteria were: age between 18 and 65 years old; meeting AD criteria according to the DSM-IV-TR (APA, 2000) and ICD-10 (WHO, 1992) or Prigerson and Jacobs' (2001) criteria for CG. Exclusion criteria were: current alcohol or drug dependence or abuse; psychosis or a severe personality disorder; a severe organic illness; malingering, showing obvious "secondary gains" (e.g., financial compensation, avoidance of work, etc.); and currently being treated in a similar treatment program. Receiving pharmacological treatment was not an exclusion criterion, but any increase and/or change in the medication during the study period implied the participant's exclusion from subsequent analyses. A decrease in pharmacological treatment was accepted. In the present study, as we have mentioned before, following ICD-10 criteria, cases of CG were considered an AD subtype, and patients with both diagnoses were included. Additionally, Prigerson and Jacobs' (2001) criteria were considered in the CG cases. Table 1 shows participants' sociodemographic and clinical data for each group and for the total sample. The type of stressful event experienced by the AD patients was: break-up relationship or divorce (n=14; 25.93%), own or family health problems (n=11; 20.37%), work/study problems (n=11; 20.37%), family problems (n=12; 21.82%), or grief (n=6; 10.91%).

2.3. Measures

2.3.1. Diagnostic

Diagnostic Interview Adjustment Disorder (Andreu-Mateu, Botella, Baños & Quero, 2008). This semi-structured interview was developed by our research team for the assessment of AD, based on data available in the literature, the diagnostic criteria from both the ICD-10 and the DSM-IV-TR, and the SCID-IV interview (First, Spitzer, Gibbon and Williams, 1999). The presence and severity of 28 symptoms related to AD are assessed on a scale from 0 to 8 (0 = Nothing at all; 8 = Very severe). This interview is currently undergoing a validation process. An optional section for evaluating Prigerson and Jacobs' (2001) criteria for CG is also included. In the present study, the number of symptoms was analyzed as an outcome measure.

2.3.2. Primary outcome measures

Inventory of Complicated Grief (ICG) (Prigerson et al. 1995). The ICG assesses the symptoms that characterize CG, and it differentiates individuals who suffer from CG symptoms, from those who are following a normal bereavement process. It consists of 19

items scored on a 5-point Likert scale ranging from 0 (“Never”) to 4 (“Always”). Scores above 25 indicate a significantly higher degree of interference in general, mental, and social functioning, worse physical health, and higher levels of physical pain. Prigerson et al. (1995) obtained high internal consistency, test-retest reliability, and convergent and criterion validity.

Adaptation of the ICG, *Inventory of Stress and Loss (ISL)* (Quero, Mor, Molés, Rachyla, Baños & Botella, submitted), to measure stress and loss symptoms derived from stressful situations. The adaptation was based on the assumption that a stressful life event always entails some kind of loss for the person. The instrument was made up of the same items included in the ICG, but the words referring to the “dead person” were substituted by words referring to the “person/situation” that had been lost due to a stressful event (e.g., loss of a job, divorce, loss of health, etc.). Two items were removed because they were not pertinent to assessing AD; therefore, the final questionnaire included 17 items. Preliminary validation data (Quero, Molés, Mor, Baños & Botella, 2014) showed excellent Cronbach coefficients in both general (0.91) and clinical AD (0.86) Spanish populations.

Beck Depression Inventory (BDI-II) (Beck Depression Inventory, Beck, Steer & Brown 1996; Spanish adaptation by Sanz, Navarro & Vázquez, 2003). The BDI-II includes 21 items, each with four possible answers (0-3), to assess severity of depression in the past 2 weeks. Spanish adaptation showed high internal consistency ($\alpha=.87$ in the general population, and $\alpha=.89$ for the clinical population) (Sanz, García-Vera, Espinosa, Fortún, & Vázquez, 2005; Sanz, Perdigón & Vázquez, 2003).

2.3.3. Secondary Outcome Measures

Posttraumatic Growth Inventory (PTGI) (Tedeschi & Calhoun, 1996). This inventory is composed of 21 items that assess the positive changes that can be experienced by people after suffering a traumatic or adverse event, using a 0-5 scale (0 = “I did not experience this change at all”; 5 = “I experienced this change to a great extent”). A global score can be calculated by adding together the scores obtained on all the responses, where high scores indicate a higher degree of perceived posttraumatic growth. High internal consistency levels have been found (Tedeschi & Calhoun 1996; So-kum Tang, 2007, Ho, Wing Chu & Yiu 2008). Preliminary data validation of the PTGI adaptation in a Spanish clinical sample with AD and CG (Molés, Quero, Nebot, Rachyla & López, 2014) showed high internal consistency (0.92) and test-retest reliability (0.95).

Maladjustment Scale (MS) (Adapted from Echeburúa, Corral & Fernández-Montalvo, 2000). This instrument assesses the level of impairment that the problem and its consequences are causing in the patient's different life areas (work, social life, leisure, partner, family, and global impairment), using 0-8 scales where 0 indicates "Not impaired" and 8 indicates "severely impaired". This scale offers good psychometric properties, and it is sensitive to the effects of treatment. In the present study, only global impairment is presented.

Clinician Severity Scale (Adapted from ADIS-IV Interview by Di Nardo, Brown & Barlow, 1994). The therapist makes a global evaluation of the patient's interference and severity on a scale from 0 ("Not impaired or without symptoms") to 8 ("Severely impaired or Very severe symptoms").

2.4. Treatment

The AD-protocol developed by Botella, Baños and Guillén (2008) was composed of 6 weekly therapy sessions lasting approximately 1.5 hours each. Two additional sessions were optional, depending on the participant's therapeutic needs. Although AD and CG are separate disorders in the ICD-11, they are included in the same category of disorders associated with stress. They have common characteristics, and so they share similar therapeutic objectives. In this regard, the main aim of the treatment applied in this study for both conditions was the reconstruction of meaning after loss, in the case of CG, and of the stressful event (e.g., break-up relationship, health problems), in the case of AD patients, in order to achieve a more positive meaning, thus allowing post-traumatic growth and learning from the negative experience. The AD-protocol includes the following therapeutic components.

1) Psychoeducation

The patient is provided with information about common reactions to a stressful event and a theoretical model that explains how the problems occur and how they are maintained. Throughout the treatment sessions, the psychoeducation component is aimed at reformulating the patient's interpretation of difficulties and his/her attitude toward problems, teaching more positive ways to manage stressful situations and introducing an alternative model explaining the onset and maintenance of AD.

2) Exposure, elaboration/processing of the stressful event

The main objective of this component is to get the person to elaborate on, find alternative meanings for, and learn from the stressful experience. This component was applied in two ways: 1) Using VR (EMMA's World). Patients' narratives of the experience are conducted within *EMMA's World* (described below), which helps participants to evoke, confront, and process the negative emotions associated with the negative event, while

remaining in a safe virtual space. 2) Traditional format. Patients' narratives of the experience are conducted within a traditional face to face therapy context.

In applying this component, instructions were given to the patient about the importance of being aware of emotions, thoughts, and behaviors without judging them. The final aim was for the participant to change the negative meaning of the stressful event into a more positive one while learning about the negative experience. In vivo exposure was also used to confront avoided situations or people related to the stressful event.

3) *Strategies from Positive Psychology*

Several strategies from Positive Psychology were included in the protocol in order to promote elaboration on, assimilation of, and learning from the experienced event. These strategies were the following: 1) "*Problem acceptance training*" (Popper, 1995). Lessons about problems as inherent to the human condition were presented to the patients. Elements include the usefulness of having problems, the significance of problem-solving, and the importance of accepting problems as endemic to the life experience. 2) "*My best virtues or strengths*" exercise (Peterson & Seligman, 2004). In this exercise, it is suggested to the patient that virtues give people the strength and capacity to cope with life. Examples of virtues are offered to the patient, including curiosity, optimism towards the future, sense of humor, etc. Patients have to analyze which ones they think they have and which ones they think it would be useful to develop and promote in order to cope with their lives. 3) "*Heuristics*" exercise. From a list of heuristics (proverbs, statements, life guidelines), patients choose the ones they find helpful in maintaining and promoting the changes achieved in their lives.

4) *Strategies from Neimeyer (2000)*

Several strategies based on Neimeyer's (2000) suggestions for CG were adapted and included in the protocol with the aim of promoting the elaboration/processing of the stressful event: 1) *The Book of Life* (adapted from Neimeyer, 2000). The aim of the book is to help the patient to remember and elaborate on meaningful events that have happened with regard to the adverse event. In writing this book, patients can use not only words, but also symbols (e.g., pictures, drawings, phrases, etc.). Through this process, patients can reflect on the negative situation, the deceased person, the broken relationship, etc. The *Book of Life* is where the patient can reexamine and sort through the jumble of memories, thoughts, and emotions that often produce confusion, pain and suffering. This process gives the person the opportunity to correct the information that was not well stored in his/her memory; that is, the person has the opportunity to rework the memories in order to create alternative meanings; 2)

the "Vital trace" exercise. Through this exercise patients try to deepen their search for positive aspects of the stressful event. According to Neimeyer (2000), people consist of the vestiges of their life experiences, both positive and negative, and all of our experiences offer opportunities to improve ourselves; 3) *Projection letter to the future*". This task requires patients to write a letter to themselves from the future. To accomplish this, they first spend 10 minutes a day for 2 to 3 days, imagining what kind of person they will be in 10 years (what they will be doing, where they will live and with whom, what their careers will be, etc.), and then they write the letter to themselves in second person (as if they were writing to a loved one) from the future. Later analysis of the letter will focus on sections that reveal the patient's ability to see "beyond" the current negative situation.

5) *Relapse prevention*

The main objective of this component is to maintain and promote therapeutic changes in the future. The therapist reviews the therapeutic objectives achieved by the patient so far, makes a summary of the main therapeutic components worked on throughout the treatment, and resolves doubts and identifies pending issues to work on in the future.

2.5. Description of EMMA's World

EMMA's World (Baños *et al.*, 2009; Botella *et al.*, 2006) was developed within a European Union project (*Engaging Media for Mental Health Applications*, IST-200-39192). It is an open and flexible adaptive VR device that adjusts its contents to the patient's needs, making possible to use this VR system to treat CG and a variety of AD cases, regardless of the stressful event that triggered them (e.g., a divorce, a firing, a serious disease, economic problems, etc.). Specifically, *EMMA's World* is used to activate and process the emotions and cognitions related to the stressful event and expose the patients to aspects they have been avoiding, in order to facilitate the emotional processing of all the aspects related to the stressful event. The underlying logic follows Foa and Kozak's approach (1986) stating that feared pathological structures need to be activated as much as possible in order to fully process the event.

In *EMMA's World*, the experienced stressful event can be "physically" represented to the patient using different symbols, such as images, 3-D objects, sounds, music, etc., that best reflect the emotions and thoughts related to the experience. Furthermore, personal elements that are significant to the patient can be introduced in the system (e.g., pictures). An important element is the *Book of Life*, a virtual book in which the patient can compile all of

the elements that help him/her to represent the most relevant moments, people or situations related to the negative experience. *EMMA*'s world also includes five different pre-defined scenarios or "landscapes": a desert, an island, a threatening forest, a snow-covered town, and meadows. These environments were designed to stimulate different emotions (relaxation, elation, sadness, etc.). Their specific use depends on the context of the session and can be selected by the therapist in real time. The therapist can also modify various aspects of these landscapes in real time (e.g., the time of day, whether it is raining or foggy, etc.) with the aim of reflecting and enhancing the emotion the user is experiencing, or inducing certain emotions. In sum, the different elements included in the system and the landscapes of *EMMA's World* are designed to help patients confront, accept, and manage the previous emotions and negative experiences in their lives, and find and develop new meanings and positive emotions, while working in a timeless place and a safe therapeutic context. More detailed descriptions of *EMMA's World* can be found elsewhere (Baños *et al.*, 2009; Baños *et al.*, 2011; Botella *et al.*, 2006; Botella, Quero *et al.*, 2006).

2.6. Technical aspects

The following devices were used: two PCs, a large screen on which the environment was projected, two projectors, a wireless pad, and a system of speakers. These devices were placed in a 5x9 meter room. PC#1 had the graphical output from its graphic card connected to two projectors (with a resolution of 1024x768 pixels and a power of 2000 lumens). They were used to project the environment on a horizontal methacrylate screen measuring 4x1.5 meters, placed in the middle of the room. The patient could see the virtual environment, interact with it, and navigate through it using the wireless pad. The therapist could control the application and the features of the virtual environment that were shown to the patient with PC#2. The sound system consisted of several speakers distributed in the room to conform to a 5.1 configuration.

2.7. Procedure

All the participants who met the inclusion criteria signed the informed consent to participate in the study. Then they underwent two assessment sessions lasting 1.5 hours each. In the first assessment session, the diagnosis of AD or CG was confirmed, and self-report measures were completed. In the second assessment session, the therapeutic goals were established. The randomization of the participants took place after the assessment sessions were conducted. Therapists and participants involved in the trial were blind to treatment allocation during the assessment. Participants were assigned to either the *EMMA* (N = 18), Traditional (N = 18), or Waiting list control (N = 18) conditions based on a computer

generated randomization list created by the “Random Allocation Software”; version 1.0. Then participants assigned to the treatment conditions were treated in 6 to 8 treatment sessions lasting approximately 1.5 hours each, with the only difference being the use of VR for the processing/elaboration of the stressful event in the EMMA condition. Participants included in the control condition did not receive any intervention during a 6-week period, after which they were assessed again and then offered the possibility of receiving one of the two treatment conditions. Once the treatment was over, all the participants were assessed again at post-treatment and at 6 and 12-month follow-ups.

Five therapists participated in the study. They had a PhD or a Master’s degree in Psychology. All of them were trained in VR techniques and CBT programs for Emotional Disorders. In addition, they had received training in this treatment protocol by senior clinicians. They were supervised by senior clinicians with PhDs in weekly sessions. Moreover, all of the assessment and treatment sessions were video-taped in order to supervise the performance of each therapist.

2.8. Data Analyses

Group differences at pre-treatment on demographic data and clinical variables were evaluated using chi-square tests for categorical variables and one-factor analyses of variance (ANOVAs) for continuous data. Intent-to-treat (ITT) mixed models analyses without any ad hoc imputation were used to account for missing data (Chakraborty & Gu, 2009). A mixed model approach is appropriate for RCTs with multiple time points and pre-to post-only designs, it does not assume that the last measurement is stable, and it is conducted using all available observations (Gueorguieva, & Krystal, 2004; Salim, Mackinnon, Christensen, & Griffiths, 2008). A linear mixed model for each outcome measure was implemented with *time* (pre and post) as within-group factor and *group* (EMMA, Traditional and WL) as between-group factor using the MIXED procedure with a random intercept for subject. Because the random effect had one level, an identity covariance structure was specified to model the covariance structure of the intercept. Separate mixed-model analyses were performed to examine changes from baseline in each intervention at posttest and 6- and 12-month follow-ups. Significant effects were followed up with pairwise contrasts. Effect sizes (Cohen’s *d*) were calculated for within- and between-group comparisons, based on Botella and Sánchez-Meca (2015). Power analysis to compute the achieved power of statistical tests for the primary outcome measures were performed using the G*Power 3.1.9.2 given α , sample size, and effect size (by Cohen’s *d*). Finally, Chi-squared tests were performed based on completers in order to examine the differential clinically significant improvement rates,

following the Jacobson and Truax (1991) indexes, for the primary outcomes scores at post-treatment and 6- and 12-month follow-ups in the two treatment conditions just considering those participants who completed the treatment protocol. Statistical analyses were performed using the IBM SPSS version 23 for Windows.

3. RESULTS

3.1. Participant flow and attrition

Participants were recruited between January 2009 and April 2011. Initially, as the flow diagram shows (see Figure 1), 97 people were interested in the study, and 67 of them were assessed for eligibility criteria. Thirteen participants were excluded from the study for several reasons (see Figure 1). Thus, 54 participants were randomly allocated to each experimental condition (EMMA condition, $N = 18$; Traditional condition, $N = 18$; and WL, $N = 18$). However, during treatment, 8 participants dropped out of the study. No significant differences in attrition rates were found at post-treatment between conditions. After treatment completion, 1 participant in the traditional condition worsened and was excluded from the study. This participant was successfully treated years ago for an obsessive-compulsive disorder that reappeared in the follow-up assessment period of the present study due to a new stressful situation in his life, and so more therapy sessions were needed. Therefore, in this condition, 14 participants completed the 6-month follow-up assessment. Finally, at 12-month follow-up, a total of 24 participants attended the assessment session: 10 participants in the traditional condition and 14 in the EMMA condition. Data were missing completely at random (MCAR) ($p > .50$).

3.2. Baseline data and participants characteristics

Baseline analyses showed no significant differences between the three groups at pre-treatment on any demographic variables, diagnostic subtype, length of the problem, medication intake variables, and outcomes measures, except for the PTGI ($F(2,51)=5.187$, $p < .05$), where the WL control group scored significantly higher than the two treatment conditions (EMMA and Traditional). No variations in the medication were produced throughout the study.

3.3. Differential effectiveness of the three experimental conditions: changes in outcome measures at pre and post-treatment

Primary outcomes

A significant time x group interaction was found for *BDI-II* ($F(2, 46.78) = 3.575, p < .05$) and *ISL/ICG* ($F(2, 44.36) = 27.898, p < .05$). Within-group comparisons showed significant reductions on both primary outcome measures for the Traditional [*BDI*: $F(1, 47.14) = 22.650, p < .001$; *ISL/ICG*: $F(1, 44.21) = 18.385, p < .001$] and EMMA [*BDI*: $F(1, 47.14) = 17.304, p < .001$; *ISL/ICG*: $F(1, 44.21) = 18.129, p < .001$], conditions, corresponding to large effect sizes (from 1.02 to 1.24), but no significant change was found in the WL group (see Table 2 and Figure 2). For the *BDI-II*, between-group comparisons revealed that post-treatment scores were significantly lower in the Traditional and EMMA groups, compared to WL, with moderate-large effect sizes (-.89 and -.78, respectively) (see Table 3). For the *ISL/ICG*, between-group comparisons revealed that participants in the Traditional condition scored significantly lower at post-treatment than the WL, with a large effect size ($d = -.92$) (see table 3 for details). The power achieved by the statistical analysis was $\beta > .80$ (ranging from .98 to .99) and α of .05 for both primary outcome measures.

Secondary outcomes

Regarding *PTGI*, pre-treatment scores on this measure were entered as a covariate to account for baseline differences. Results indicated a significant time x group interaction effect ($F(2, 44.862) = 13.733, p < .001$). Within-group comparisons revealed a significant pre-to-post change in the two treatment groups ($F(1, 45.112) = 49.962, p < .001$, for both groups), with large effect sizes (from -1.62 to -2.05), and no significant change in the WL (table 2). At post-treatment, no statistically significant between-group differences were found (Table 3).

For both the *Maladjustment Scale* and *Clinician Severity Scale*, a significant time x group interaction was found [$F(2, 48.32) = 5.911, p < .01$] ($F(2, 25.617) = 24.088, p < .001$), respectively]. Within-group comparisons revealed significant reductions for both treatments [*Maladjustment Scale*, Traditional: $F(1, 49.19) = 33.981, p < .001$; EMMA: $F(1, 47.89) = 21.426, p < .001$] (*Clinician Severity Scale*, Traditional: $F(1, 26.51) = 29.997, p < .001$; EMMA: $F(1, 24.220) = 93.938, p < .001$], corresponding to large effect sizes, but no significant change was found in WL (see Table 2). Between-group comparisons showed that post-treatment scores on these measures were significantly lower in both treatments compared to WL, achieving large effect sizes (from -.81 to -1.94) (Table 3).

3.4. Differential effectiveness of the two treatment conditions: changes in outcome measures at pre-, 6-, and 12-month follow-ups

Separate mixed model analysis for each primary and secondary outcome measure yielded no significant time x group interaction effect. Within-group comparisons revealed a significant change (from pre- to 6- and 12-month follow-ups, respectively) in both treatments for all primary and secondary outcome measures (all $ps < .001$), corresponding to large effect sizes (see table 2 for details). No statistically significant differences between the two treatments (EMMA vs. Traditional) were found at the 6-month or 12-month follow-up (Table 3). However, larger effect sizes were obtained in the EMMA condition at the 12-month follow-up, compared to the Traditional condition (see Table 2 and Figure 2).

3.5. Clinically meaningful improvement

Table 4 shows the percentages of participants in each condition who were *recovered*, *improved*, had *no change*, or were *impaired*, according to the ISL/ICG and BDI measures, and based on the classification proposed by Iraurgi (2010) and Kupfer (1991). In general, better results were obtained for the *primary outcome measure* most closely related to the problem, the ISL/ICG. At post-treatment, more participants (who had completed the treatment) in both conditions were included in the categories recovered or improved. Then, the percentage of *recovered* improved slightly at the 6-month follow-up in both conditions, but in the long term, more participants in the EMMA condition were included in the recovered category, whereas a decrease was found in the Traditional condition. In the case of the BDI-II, again significantly more participants were included in the recovered category in the EMMA condition than in the Traditional condition.

4. DISCUSSION

The main objective of the present study was to test, in a controlled study, the long-term efficacy of a CBT protocol for AD and CG in a traditional format versus a format supported by VR (EMMA's World), and compared to a WL control group. First, data showed that both treatment conditions resulted in statistically significant improvements with large effects sizes on both primary and secondary outcome measures, and this improvement did not occur in the WL control condition. Therefore, the first hypothesis was confirmed. These data agree with those obtained in previous studies (Andreu-Mateu *et al.*, 2012; Baños *et al.*, 2011), and they confirm the usefulness of this psychological treatment protocol supported by VR for AD and CG. These results are also consistent with the conclusions reported by review studies

on the efficacy of VR exposure therapy for the treatment of anxiety disorders: systematic reviews (Gregg & Tarrier, 2007; McCann et al., 2014; Meyerbroeker & Emmelkamp, 2010) and meta-analyses (Morina et al., 2015; Oprış *et al.*, 2012; Parsons & Rizzo, 2008; Powers & Emmelkamp, 2008), and by meta-analytic studies specifically addressing the use of VR for stress-related disorders (Botella, Serrano, Baños & Garcia-Palacios, 2015; Gonçalves, Pedrozo, Coutinho, Figueira & Ventura, 2012; Motraghi, Seim, Meyer & Morissete, 2014). Finally, our results are also consistent with those obtained in studies on the traditional treatment for AD (Andrews, 1995; Araoz & Carrese, 1996; Benton & Lynch, 2005, Strain, 1995; van der Klink, *et al.*, 2003) and CG (Neimeyer, 2000).

Second, as expected, no statistically significant differences were found between the two treatment groups at any assessment moment. However, at the 12-month follow-up, additional improvements were observed in the EMMA condition, with larger effect sizes in the long term, compared to the Traditional condition (see Table 2). Similarly, results for clinically significant change estimations indicated an advantage for the EMMA condition on the primary outcome measures (ISL/ICG and BDI), with the percentage of recovered participants being significantly higher at 12-month follow-up in this condition, compared to the traditional one. In any case, it is important to point out that scores on the BDI-II were not very high at pre-treatment. Therefore, the second hypothesis was partially confirmed.

Finally, as hypothesized, significant improvements achieved in both treatments for all primary and secondary outcome measures were maintained in the medium (6-month) and long term (12-month).

Based on the results obtained in the present study, we can conclude that both treatment conditions were effective at post-treatment, but the EMMA condition was slightly superior on the outcome measures and the clinically meaningful change in the long term. Furthermore, results obtained from 18 patients included in this study (7 from the traditional condition and 11 from the EMMA condition) after treatment, showed that 66.7% of them preferred the VR condition, whereas only 33.6% preferred the traditional one. Our data indicate a preference for the EMMA system, but this result cannot be generalized to all patients since the sample size was small. Moreover, it should be kept in mind that these patients were willing to receive psychological treatment supported by VR, and not all patients would necessarily have a positive opinion about technologies. Additionally, most of the participants (83.3% in EMMA condition and 72.2% in the Traditional Condition) held a university degree which might also explain why they preferred VR and, this might not be representative for the general population.

On the other hand, in the case of these problems, the positive results obtained in the long term are especially relevant, considering the few studies available in the literature about the maintenance of therapeutic gains. Results from this study offer additional support for the positive and lasting effects of VR in the psychological treatment field (e.g., Botella *et al.*, 2007; Emmelkamp *et al.*, 2002; Rothbaum, Hodges, Anderson, Price & Smith, 2002; Ready, Pollack, Rothbaum & Alarcón, 2006). Other advantages of the VR environment used in the present study should be pointed out. First, as mentioned in the introduction, EMMA's World adapts its contents in relation to the affective manifestations the users express at any time during the therapy session, resulting in the creation of a personalized and clinically meaningful environment, which, in turn, favors the complete immersion in the narrative and the elaboration and assimilation of the events. Second, the system uses symbols, landscapes and other personalized elements to create a context that "represents or symbolizes" the situation. The use of VR and symbols to conduct exposure/elaboration provides a safe context where patients can experience and confront their problems without the inconvenience of real life. In fact, patients perceive VR exposure to be less aversive than the traditional exposure technique (Baños *et al.*, 2009; Botella *et al.*, 2016).

In summary, this study follows along the lines of existing literature supporting the use of technology (in this case VR) in the field of psychological treatments (Freeman *et al.* 2017) because, as the data point out, information and communication technologies can improve psychological treatments by providing a cost-effective way to deliver empirically validated treatments for a variety of psychological problems (Newman *et al.*, 2011), in this case AD. Undoubtedly, this work contributes to achieving the crucial revolution anticipated through the use of VR in the mental health field (Freeman *et al.*, 2017; Linder *et al.*, 2017). Indeed, studies like the present one indicate that this revolution is already underway.

However, the present study has some limitations that should be highlighted. First, the sample size was small and most of the participants had university degree, limiting the generalization of the results. The level of education seems to be well above average, which might also limit the generalizability of the results. On the other hand, as mentioned before, for the diagnosis DSM-IV-TR and ICD-10 criteria were followed in the present study, and the diagnostic interview for AD is currently under validation process. The Diagnostic Interview for Adjustment Disorder (DIAD) (Comelius *et al.*, 2014) could not be used because it was not available at the time the study was conducted. In addition, it would have been desirable to include an anxiety measure as a primary outcome. Finally, all participants received the complete AD-protocol, and so it is not possible to determine the specific

contribution of each therapeutic component to the patients' improvement. Related to this, future studies should consider evaluating whether AD and CG patients differ in their response to this treatment protocol, in order to detect the need to tailor the treatment to each stress-related disorder included in the present study.

A future guideline in the field of using technology to improve psychological interventions in AD and CG is the use of the Internet to complement the therapy. Preliminary data are already available on the usefulness of an online emotion regulation system (TEO) to apply homework assignments over the Internet in treating AD (Quero, Molés, Pérez-Ara, Botella & Baños, 2012). This system allows patients to practice the elaboration/processing of the stressful event at home during the period between sessions. A system such as TEO offers great adaptability and flexibility in adjusting the therapeutic homework to each patient's characteristics and needs, focusing on the specific significance of his/her problem.

To conclude, the results obtained in the present study have contributed to increasing the knowledge and available resources to deal with AD and CG. It is the first controlled study to compare an AD-protocol applied face-to-face to the same protocol supported by VR and a waiting list control group in the treatment of these stress-related disorders. The possibility of having an effective treatment protocol is particularly relevant in the AD field, considering the few controlled studies found in the literature about AD treatment, in addition to its high prevalence (Carta *et al.*, 2009), the impairment in social and work functioning associated with this problem (Van der Klink *et al.*, 2003), the potential AD has of presenting life-threatening features (Casey *et al.*, 2015), and in the long term, the burden for the individual, his/her work place, society, and the economy (Lagerveld *et al.*, 2012).

Acknowledgments

The research presented in this paper was funded in part by Plan 2011 de Promoción de la Investigación UJI-Proyectos BANCAIXA (P11B2011-20), Ministerio de Economía y Competitividad (Spain) (Plan Nacional I+D+I. PSI2013-41783-R), and CIBER. CIBER Fisiopatología de la Obesidad y Nutrición is an initiative of ISCIII.

References

- American Psychiatric Association (2000). Diagnostic and statistical manual for mental disorders (4th edition, revised text) (DSM-IV-TR). Washington: DC.
- Andreu-Mateu, S., Botella, C., Quero, S., Guillén, V., & Baños, R. M. (2012). El tratamiento de los trastornos adaptativos: cuando el estímulo estresante sigue presente. *Behavioral Psychology/Psicología Conductual*, 20, 323-348.
- Andrews, A. (1985). Treatment outlines for the management of anxiety status: The quality assurance project. *Australian and New Zealand Journal of Psychiatry*, 19, 138-151.
- Araoz, D. L., & Carrese, M. A. (1996). Solution-oriented brief therapy for adjustment disorders: A guide for providers under managed care. New York, N.Y: Brunner/Mazel.
- Baños, R. M., Guillén, V., Quero, S., García-Palacios, A., & Botella, C. (2011). A virtual reality system for the treatment of stress-related disorders: A preliminary analysis of efficacy compared to a standard cognitive. *International Journal of Human-Computer Studies*, 69, 602-613.
- Baños, R. M., Botella, C., Guillén, V., García-Palacios, A., Quero, S., Bretón-López, J., & Alcañiz, M. (2009). An adaptive display to treat stress-related disorders: The EMMA's world. *British Journal of Guidance and Counseling*, 37, 347-356.
- Beck, A. T., Steer, R. A., & Brown, G. K. (1996). Manual for the Beck Depression Inventory-II. TX Psychological Corp, San Antonio.
- Ben-Itzhak, S., Bluvstein, I., Schreiber, S., Aharonov-Zaig, I., Maor, M., Lipnik, R., & Bloch, M. (2012). The effectiveness of brief versus intermediate duration psychodynamic psychotherapy in the treatment of adjustment disorder. *Journal of Contemporary Psychotherapy*, 42, 249-256.
- Benton, T. D., & Lynch, J., 2005. EMedicine. Adjustment disorders. <http://www.emedicine.com/Med/topic3348.htm>. (accessed 07.06.10).
- Boelen, P.A., de Keijser, J., van den Hout, M. & van den Bout, J. (2007). Treatment of complicated grief: A comparison between cognitive-behavioral therapy and supportive counseling. *Journal of Clinical and Consulting Psychology*, 75, 277-284.
- Boelen, P. A., & Smid, G. E. (2017). Disturbed grief: Prolonged grief disorder and persistent complex bereavement disorder. *The BMJ*, 357, j2016
- Botella, C., Baños, R. M., & Guillén, V. (2008). Creciendo en la adversidad. Una propuesta de tratamiento para los trastornos adaptativos. In C. Vázquez & G. Hervás (Eds.), *Psicología Positiva Aplicada* (pp. 129-154). Bilbao: DDB.

- Botella, C., Baños, R. M., Rey, B., Alcañiz, M., Guillén, V., Quero, S., & García-Palacios, A. (2006). Using an Adaptive Display for the Treatment of Emotional Disorders: A preliminary analysis of effectiveness. *Work-In-Progress, CHI*. Montréal, Québec, Canada. pp. 586-591.
- Botella, C., García-Palacios, A., Villa, H., Baños, R. M., Quero, S., Alcañiz, M., & Riva, G. (2007). Virtual reality exposure in the treatment of panic disorder and agoraphobia: A controlled study. *Clinical Psychology and Psychotherapy*, 14, 164-175.
- Botella, C., Osma, J., García Palacios, A., Guillén, V., & Baños, R. M. (2008). Treatment of complicated grief using virtual reality. A case report. *Death Studies*, 32, 674-692.
- Botella, C., Pérez-Ara, M. A., Bretón-López, J., Quero, S., García Palacios, A. & Baños, R. M. (2016). In vivo versus augmented reality exposure in the treatment of small animal phobia: a randomized controlled trial. *Plos One* 11(2): e0148237.
- Botella, C., Quero, S., Lasso de la Vega, N., Baños, R. M., Guillén, V., García Palacios, A., & Castilla, D. (2006). Clinical issues in the application of virtual reality to treatment of PTSD. In M. Roy (Ed.), *Novel Approaches to the Diagnosis and Treatment of Posttraumatic Stress Disorder* (pp. 183-195). Amsterdam: IOS Press.
- Botella, J., & Sanchez-Meca, J. (2015). *Meta-análisis en Ciencias Sociales y de la Salud*. Madrid, Spain: Síntesis
- Botella, C., Serrano, B., Baños, R., & García-Palacios, A. (2015). Virtual reality exposure-based therapy for the treatment of posttraumatic stress disorder. A review of its efficacy, the adequacy of the treatment protocol, and its acceptability. *Neuropsychiatric Disease and Treatment*, 11, 2533-45. doi: 10.2147/NDT.S89542.
- Bryant, R. A., Kenny, L., Joscelyne, A., Rawson, N., Maccallum, F., Cahill, C., ... & Nickerson, A. (2014). Treating prolonged grief disorder: A randomized clinical trial. *JAMA psychiatry*, 71(12), 1332-1339.
- Carta, M. G., Balestrieri, M., Murru, A., & Carolina-Hardoy, M. C. (2009). Adjustment Disorder: epidemiology, diagnosis and treatment. *Clinical Practice and Epidemiology in Mental Health*, 5, 1-15.
- Casey, P. (2009). Adjustment Disorder: epidemiology, diagnosis and treatment. *CNS Drugs*, 23, 927-938.
- Casey, P. (2014). Adjustment Disorders: New developments. *Current Psychiatry Reports*, 16, (6), 451-459.
- Casey, P., Jabbar, F., O'Leary, E., & Doherty, A. M. (2015). Suicidal behaviours in adjustment disorder and depressive disorder. *Journal of Affective Disorders*, 174, 441-446.

- Chakraborty, H., & Gu, H. (2009). *A mixed model approach for intent-to-treat analysis in longitudinal clinical trials with missing values* (Report No. MR-0009-0903). Research Triangle Park, NC: RTI Press.
- Comelius, L. R., Brouwer, S., de Boer, M. R. *et al* (2014). Development and validation of the Diagnostic Interview of Adjustment Disorder (DIAD). *International Journal of Methods in Psychiatric Research*, 23 (2), 192-207
- Crunk, A.E., Burke, L.A. & Robinson III, E.H.M. (2017). Complicated grief: An evolving theoretical landscape. *Journal of Counseling & Development*, 95, 226-233.
- Currier, J. M., Holland, J. M., & Neimeyer, R. A. (2010). Do CBT-Based Interventions Alleviate Distress Following Bereavement? A Review of the Current Evidence. *International Journal of Cognitive Therapy*, 3, 77-93.
- Cvetek, R. (2008). EMDR treatment of distressful experiences that fail to meet the criteria for PTSD. *Journal of EMDR Practice and Research*, 2 (1), 2-14.
- Di Nardo, P. A., Brown, T. A., & Barlow, D. H. (1994). Anxiety disorders interview schedule for DSM-IV: Lifetime Version (ADIS-IV), San Antonio, Texas.
- Echeburúa, E., Corral, P., & Fernández-Montalvo, J. (2000). Escala de inadaptación: propiedades psicométricas en contextos clínicos. *Análisis y Modificación de Conducta*, 26, 325-340.
- Emmelkamp, P. M. G., Krijn, M., Hulsbosch, A. M., de Vries, S., Schuemie, M. J., & van der Mast, C. A. P. G. (2002). Virtual reality treatment versus exposure in vivo: a comparative evaluation in acrophobia. *Behaviour Research and Therapy*, 40, 509-516.
- Foa, E. B., & Kozak, M. J. (1986). Emotional processing of fear: Exposure to corrective information. *Psychological Bulletin*, 99, 20-35.
- Freeman, D., Reeve, S., Robinson, A., Ehlers, A., Clark, D., Spanlang, B. & Slater, M. (2017). Virtual reality in the assessment, understanding, and treatment of mental health disorders. *Psychological Medicine*. Doi: 10.1017/S003329171700040X
- Goncalves, R., Pedrozo, A. L., Coutinho, E. S. F., Figueira, I., & Ventura, P. (2012). Efficacy of Virtual Reality Exposure Therapy in the Treatment of PTSD: A Systematic Review. *Plos One*, 7, 12. e48469.
- Gonzalez-Jaimes, E. I., & Turnbull-Plaza, B. (2003). Selection of psychotherapeutic treatment for adjustment disorder with depressive mood due to acute myocardial infarction. *Archives of Medical Research*, 34, 298-304.
- Gregg, L., & Tarrier, N. (2007). Virtual reality in mental health: a review of the literature. *Social Psychiatry and Psychiatric Epidemiology*, 42, 343-354.

- Gueorguieva, R., & Krystal, J. H. (2004). Move Over ANOVA. *Archives of General Psychiatry*, 61(3), 310. doi:10.1001/archpsyc.61.3.310
- Ho, S. M., Chu, K. W., & Yiu, J. (2008). The relationship between explanatory style and posttraumatic growth after bereavement in a non-clinical sample. *Death Studies*, 32, 461-478.
- Hsiao, F. H., Lai, Y. M., Chen, Y. T., Yang, T. T., Liao, S. C., Ho, R. T. H., Ng, S. M., Chan, C. L., & Jow, G. M. (2014). Efficacy of psychotherapy on diurnal cortisol patterns and suicidal ideation in adjustment disorder with depressed mood. *General Hospital Psychiatry*, 36, 214-219.
- Jacobson, N. S., & Truax, P. (1991). Clinical significance: a statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology*, 59, 12-19.
- Kryzhanovskaya, L., & Canterbury, R. (2001). Suicidal behavior in patients with adjustment disorders. *Crisis*, 22, 125-131.
- Latham, A. E. y Prigerson, H. G. (2004). *Suicide and Life-Threatening Behavior*, 34 (4), 350-362.
- Lagerveld, S., Blonk, R., Brenninkmeijer, V., Wijngaards-de Meij, L. & Schaufeli, W. (2012). Work-Focused Treatment of Common Mental Disorders and Return to Work: A Comparative Outcome Study. *Journal of Occupational and Health Psychology*, 17, 220-234.
- Li, J., Laursen, T. M., Precht, D. H., Olsen, J., & Mortensen, P. B. (2005). Hospitalization for mental illness among parents after the death of a child. *The New England Journal of Medicine*, 352, 1190-6. doi: 10.1056/NEJMoa033160
- Lindner, P., Miloff, A., Hamilton, W., Reuterskiöld, L., Andersson, G., Powers, M. & Carlbring, P. (2017). Creating state of the art, next-generation Virtual Reality exposure therapies for anxiety disorders using consumer hardware platforms: Design considerations and future direction. *Cognitive Behaviour Therapy*, 46, 404-420. doi: 10.1080/16506073.2017.1280843
- Lundorff, M., Holmgren, H., Zachariae, R., Farver-Vestergaard, I. & M. O'Connor (2017). Prevalence of prolonged grief disorder in adult bereavement: A systematic review and meta-analysis. *Journal of Affective Disorders*, 212, 138-149.
- Maercker, A., Brewin, C. R., Bryant, R. A., Cloitre, M., van, Ommeren., Jones, L. M., Humayan, A., Kagee, A., Llosa, A. E., Rousseau, C., Somasundaram, D. J., Souza, R., Suzuki, Y., Weissbecker, I., Wessely, S. C., First, M.B., & Reed, G. M. (2013). Diagnosis and classification of disorders specifically associated with stress: proposals for ICD-11. *World. Psychiatry*, 12, 198-206.
- McCann, R. A., Armstrong, C. M., Skopp, N. A., Edwards-Stewart, A., Smolenski, D. J., June, J. D., Metzger-Abamukong, M., & Reger, G. M. (2014). Virtual reality exposure therapy for the

- treatment of anxiety disorders: An evaluation of research quality. *Journal of Anxiety Disorders*, 28, 625-631.
- Meyerbröker, K., & Emmelkamp, P. M. G. (2010). Virtual reality exposure therapy in anxiety disorders: a systematic review of process-and-outcome studies. *Depression and Anxiety*, 27, 933-944.
- Molés, M., Quero, S., Nebot, S., Rachyla, I., & López, M. T. (2014). Datos preliminares de validación del Inventario de Crecimiento Postraumático en población clínica española con trastornos relacionados con el estrés. Paper presented at II congreso nacional de psicología positiva. May, Oropesa (Spain).
- Morina, N., Ijntema, H., Meyerbröcker, K. & Emmelkamp, P.M.G. (2015). Can virtual reality exposure therapy gains be generalized to real-life? A meta-analysis of studies applying behavioral assessments. *Behaviour Research and Therapy*, 74, 18-4.
- Motraghi, T. E., Seim, R. W., Meyer, E. C., & Morissette, S. B. (2014). Virtual reality exposure therapy for the treatment of posttraumatic stress disorder: A methodological review using CONSORT guidelines. *Journal of Clinical Psychology*, 70, 197-208.
- Neimeyer, R. A., & Currier, J. M. (2009). Grief Therapy: Evidence of Efficacy and Emerging Directions. *Current Directions in Psychological Science*, 18, 352-356.
- Neimeyer, R. A. (2000). *Lessons of Loss*. Florida: Keystone Heights.
- Newman, M.G., Szkodny, L.E., Llera, S. J. & Przeworski, A. (2011). A review of technology-assisted self-help and minimal contact therapies for anxiety and depression: Is human contact necessary for therapeutic efficacy?. *Clinical Psychology Review*, 31, 89-103.
- Opriş, D., Pinte, S., García-Palacios, A., Botella, C., Szamosközi, S., & David, D. (2012). Virtual reality exposure therapy in anxiety disorders: a quantitative meta-analysis. *Depression and Anxiety*, 29, 85-93.
- Parsons, T. D., & Rizzo, A. A. (2008). Affective outcomes of virtual reality exposure therapy for anxiety and specific phobias: A meta-analysis. *Journal of Behavior Therapy and Experimental Psychiatry*, 39, 250-261.
- Pelkonen, M., Marttunen, M., Henriksson, M., & Lönnqvist, J. (2007). Adolescent adjustment disorder: Precipitant stressors and distress symptoms of 89 outpatients. *European Psychiatry*, 22, 288-295.
- Peterson, C., & Seligman, M. E. (2004). *Character strengths and virtues: A handbook and classification*. Washington, DC: American Psychological Association.
- Popper, K. (1995). *La responsabilidad de vivir*. Barcelona: Paidós.

- Powers, M. B., & Emmelkamp, P. M. (2008). Virtual reality exposure therapy for anxiety disorders: A meta-analysis. *Journal of Anxiety Disorders*, 22, 561- 569.
- Prigerson, H. G., Bierhals, A. J., Kasl, S. V., Reynolds, C. F., Shear, M. K., Day, N., ...& Jacobs, S. (1997). Traumatic grief as a risk factor for mental and physical morbidity. *American Journal of Psychiatry*, 154, 616-623.
- Prigerson, H. G., Maciejewski, P. K., Reynolds, C. F., Bierhals, A. J., Newsom, J. T., Fasiczka, A., Frank, E., Doman, J., & Miller, M. (1995). Inventory of Complicated Grief: A scale to measure maladaptive symptoms of loss. *Psychiatry Research*, 59, 65-79.
- Prigerson, H., & Jacobs, S. (2001). Traumatic grief as a distinct disorder: A rationale, consensus criteria, and a preliminary empirical test. In M. Stroebe, W. Stroebe, R. O. Hansson, & H. Schut (Eds.), *Handbook of bereavement research: Consequences, coping, and care*. (pp. 613-645). Washington, DC: American Psychological Association Press.
- Quero, S., Molés, M., Mor, S., Baños R., & Botella, C. (2014). Spanish Validation of the Complicated Grief Inventory adapted for Adjustment Disorders. Paper presented at *ABCT's 48th Annual Convention*. November, Philadelphia (USA).
- Quero, S., Molés, M., Pérez, M. A., Botella, C., & Baños, R. M. (2012). An online emotional system to deliver homework assignments for treating adjustment disorders. In B. K. Wiederhold & G. Riva (Eds.), *Annual review of cybertherapy and telemedicine* (pp. 273-278). Interactive Media Institute, Amsterdam: IOS Press. ISBN: 978-1-61499-120-5.
- Ready, D. J., Pollack, S., Rothbaum, B. O., & Alarcon, R. D. (2006). Virtual reality exposure for veterans with posttraumatic stress disorder. *Journal of Aggression, Maltreatment and Trauma*, 12, 199-220.
- Rothbaum, B. O., Hodges, L., Anderson, P. L., Price, L., & Smith, S. (2002). Twelve-month follow-up of virtual reality and standard exposure therapies for the fear of flying. *Journal of Consulting and Clinical Psychology*, 70, 428-432.
- Salim, A., Mackinnon, A., Christensen, H., & Griffiths, K. (2008). Comparison of data analysis strategies for intent-to-treat analysis in pre-test–post-test designs with substantial dropout rates. *Psychiatry Research*, 160(3), 335–345. doi:10.1016/j.psychres.2007.08.005
- Sanz, J., García-Vera, M-P., Espinosa, R., Fortún, M., & Vázquez, C. (2005). Adaptación española del inventario para la depresión de Beck- II (BDI-II): Propiedades psicométricas en pacientes con trastornos psicológicos. *Clínica y Salud*, 16, 121-142.

- Sanz, J., Navarro, M. E., & Vázquez, C. (2003). Adaptación española del Inventario para la Depresión de Beck-II (BDI-II): Propiedades psicométricas en estudiantes universitarios. *Análisis y Modificación de Conducta*, 29, 239-288.
- Sanz, J., Perdigón, A., & Vázquez, C. (2003). Adaptación española del Inventario para la Depresión de Beck-II (BDI-II): 2. Propiedades psicométricas en población general. *Clínica y Salud*, 14, 249-280.
- Shear, M.K. (2015). Complicated Grief. *The New England Journal of Medicine*, 372, 153-160.
- Shear, M.K., Frank, E., Houck, P.R. & Reynolds, C.F. (2005). Treatment of complicated grief: A randomized controlled trial. *Journal of the American Medical Association*, 293, 2601-2608.
- Shear, M.K., Wang, Y., Skritskaya, M., Duan, N., Mauro, C. & Ghesquiere, A. (2014) Treatment of complicated grief in elderly persons: a randomized clinical trial. *Journal of the American Medical Association*, 71, 1287-1295.
- Simon, N. M., Shear, K. M., Thompson, E. H., Zalta, A. K., Perlman, C., Reynolds, C. F. ... & Silowash, R. (2007). The prevalence and correlates of psychiatric comorbidity in individuals with complicated grief. *Comprehensive Psychiatry*, 48(5), 395-399.
- So-Kum Tang, C. S. (2007). Posttraumatic growth of Southeast Asian survivors with physical injuries: Six months after the 2004 Southeast Asian earthquake-tsunami. *Australasian Journal of Disaster and Trauma Studies*, 1 ISSN: 1174-4707.
- Srivastava, M., Talukdar, U., & Lahan, V. (2011). Meditation for the management of adjustment disorder anxiety and depression. *Complementary Therapies in Clinical Practices*. 17, 241-245.
- Steinhardt, M., & Dolbier, C. (2008). Evaluation of a resilience intervention to enhance coping strategies and protective factors and decrease symptomatology. *Journal of American College Health*, 56, 445-453.
- Strain, J. J. (1995). Adjustment disorders. In G. O. Gabbard (Ed.). *Treatments of psychiatric Disorders*, second ed. (pp. 1656-1665). Washington, DC: APA Press.
- Stroebe, M. S., Schut, H. A. W., & Stroebe, W. (2007). Health outcomes of bereavement. *Lancet*, 370, 1960-73. doi: 10.1016/S0140-6736(07)61816-9.
- Tedeschi, R. G., & Calhoun, L. G. (1996). The Posttraumatic Growth Inventory: measuring the positive legacy of trauma. *Journal of Traumatic Stress*, 9, 455-471.
- Van der Klink, J. L., Blonk, R. W., Schene, A. H. & van Dijk, F. J. (2003). Reducing long term sickness absence by an activating intervention in adjustment disorders. *Occupational and Environmental Medicine*, 60, 429-437.

World Health Organization (1992). International Classification of Diseases (ICD-10) (10th Edition).
Madrid: Meditor.

Yaseen, Y. A. (2017). Adjustment disorder: Prevalence, sociodemographic risk factors, and its subtypes in outpatient psychiatric clinic. *Asian journal of psychiatry*, 28, 82-85.

Accepted Article

Table 1. Participants' descriptive data about demographic variables, and diagnosis variables.

	EMMA (N = 18)	TRAD (N = 18)	WL (N = 18)	Total (N = 54)
Age				
Range	21-49	18-42	21-50	18-50
Mean (SD)	34.67 (8.63)	30.06 (7.83)	34.67 (8.35)	33.13 (8.41)
Sex				
Men	4 (22.2%)	7 (38.9%)	3 (16.7%)	14 (25.9%)
Woman	14 (77.8%)	11 (61.1%)	15 (83.3%)	40 (74.1%)
Marital status				
Single	7 (38.9%)	11 (61.1%)	8 (44.4%)	26 (48.1%)
Married/ partnered	7 (38.9%)	7 (38.9%)	7 (38.9%)	21 (38.9%)
Divorced/separated/widower	4 (22.2%)	0 (0%)	3 (16.7%)	7 (13%)
Level of studies				
Elementary/High school	3 (16.7%)	5 (27.8%)	6 (33.3%)	14 (25.9%)
University degree	15 (83.3%)	13 (72.2%)	12 (66.7%)	40 (74.1%)
Diagnosis				
AD subtype				
With depressed mood	2 (11.1%)	3 (16.7%)	2 (11.1%)	7 (13%)
With anxiety	0 (0%)	1 (5.6%)	1 (5.6%)	2 (3.7%)
With mixed anxiety and depressed mood	15 (83.3%)	12 (66.7%)	11 (61.1%)	38 (70.4%)
With disturbance of conduct	0 (0%)	0 (0%)	1 (5.6%)	1 (1.9%)
CG	1 (5.6%)	2 (11.1%)	3 (16.7%)	6 (11.1%)
Length of the problem				
Range (months to years)	3 to 20	3 to 17	5 to 10	3 to 20
Years mean (SD)	4.37 (5.29)	3.11 (4.02)	3.51 (3.38)	3.66 (4.26)
Medication (anxiolytics and/or antidepressants)				
Yes	26.7%	6.7%	31.3%	21.7%
No	73.3%	93.3%	68.8%	78.3%

Note: SD = Standard Deviation; EMMA = EMMA Intervention group; TRAD= Traditional Intervention group; WL = Waiting list control group; AD = Adjustment disorder; CG = Complicated grief.

Table 2. Means, standard deviations and within-group comparisons effect sizes for primary and secondary outcome measures.

		Pre (n=54)	Post (n=46)	6-month FW (n= 29)	12-month FW (n= 24)	Pre vs. Post	Pre vs. 6-month FW	Pre vs. 12 month FW
		Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	<i>d</i> (95% CI)	<i>d</i> (95% CI)	<i>d</i> (95% CI)
<i>ISL/ICG</i>								
	TRAD	35.72 (16.88)	17.67 (7.10)	13.71 (9.49)	11.00 (7.56)	<i>d</i> =1.02 (.55, 1.50)	<i>d</i> =1.25 (.59, 1.90)	<i>d</i> =1.40 (.59, 2.21)
	EMMA	36.33 (13.05)	19.40 (13.87)	13.53 (12.63)	5.86 (5.38)	<i>d</i> =1.24 (.48, 2.00)	<i>d</i> =1.67 (.81, 2.52)	<i>d</i> =2.23 (1.08, 3.38)
	WL	31.25 (15.93)	28.53 (14.53)	–	–	<i>d</i> =.16 (-.25, .58)	–	–
<i>BDI-II</i>								
	TRAD	23.00 (10.48)	10.40 (7.71)	8.93 (9.55)	8.70 (9.43)	<i>d</i> =1.15 (.40, 1.89)	<i>d</i> =1.28 (.57, 1.99)	<i>d</i> =1.30 (.58, 2.02)
	EMMA	23.00 (9.41)	11.87 (6.77)	7.93 (6.41)	4.21 (7.37)	<i>d</i> =1.13 (.40, 1.86)	<i>d</i> =1.53 (.80, 2.26)	<i>d</i> =1.91 (.93, 2.89)
	WL	23.17 (11.48)	19.88 (12.40)	–	–	<i>d</i> =.16 (-.09, .40)	–	–
<i>PTGI</i>								
	TRAD	30.72 (15.39)	63.67 (18.88)	58.64 (19.26)	61.30 (16.95)	<i>d</i> =-2.05 (-3.01, -.08)	<i>d</i> =-1.73 (-2.59, -.87)	<i>d</i> = -1.57 (-2.42, -.72)
	EMMA	29.50 (18.61)	61.13 (24.10)	56.00 (21.71)	63.29 (19.19)	<i>d</i> =-1.62 (-2.33, -.92)	<i>d</i> =-1.36 (-2.02, -.70)	<i>d</i> =-1.73 (-2.53, -.94)
	WL	49.28 (26.37)	51.75 (29.73)	–	–	<i>d</i> =-.09 (-.30, .12)	–	–
<i>Impairment</i>								
	TRAD	5.78 (1.00)	3.21 (1.81)	2.36 (1.34)	2.00 (1.41)	<i>d</i> =2.45 (1.44, 3.46)	<i>d</i> =3.27 (1.92, 4.62)	<i>d</i> =3.61 (2.05, 5.18)
	EMMA	5.83 (1.15)	3.80 (.86)	2.00 (1.73)	1.43 (1.28)	<i>d</i> =1.69 (.71, 2.66)	<i>d</i> =3.18 (1.79, 4.57)	<i>d</i> =3.65 (2.19, 5.12)
	WL	5.83 (1.38)	5.33 (2.44)	–	–	<i>d</i> =.35 (-.07, .77)	–	–
<i>Severity</i>								
	TRAD	5.41 (1.12)	3.17 (.75)	2.33 (1.53)	1.33 (.578)	<i>d</i> =1.91 (1.01, 2.81)	<i>d</i> =2.63 (1.43, 3.82)	<i>d</i> =3.48 (2.05, 4.91)
	EMMA	5.59 (.62)	2.70 (1.06)	2.33 (.58)	1.25 (.50)	<i>d</i> =4.45 (2.79, 6.11)	<i>d</i> =5.02 (3.12, 6.93)	<i>d</i> =6.69 (4.18, 9.20)
	WL	4.82 (1.51)	5.29 (1.50)	–	–	<i>d</i> =-.30 (-.55, -.04)	–	–

Note: Mean and standard deviations (SD) are represented. Pre= Pre-treatment. Post= Post-treatment. FW= Follow-up. *d* = Cohen's *d*. CI= Confidence Interval. EMMA = EMMA Intervention group; TRAD= Traditional Intervention group; WL = Waiting list control group; ISL= Inventory of Stress and Loss; ICG= Inventory of Complicated Grief; BDI-II= Beck Depression Inventory II; PTGI= Posttraumatic Growth Inventory; Impairment= Maladjustment Scale; Severity= Clinician Severity Scale.

Table 3. Between-group comparisons at post-treatment and 6- and 12-month follow-Up.

		Post-treatment		6-month follow-up		12-month follow-up	
		Mean diff.	<i>d</i> (95% CI)	Mean diff.	<i>d</i> (95% CI)	Mean diff.	<i>d</i> (95% CI)
<i>SLI/ICG</i>	TRAD vs. WL	-10.38*	<i>d</i> = -.92 (-1.68, -.17)	–	–	–	–
	EMMA vs. WL	-9.65	<i>d</i> = -.63 (-1.36, .11)	–	–	–	–
	EMMA vs. TRAD	.74	<i>d</i> = .15 (-.56, .87)	-.96	<i>d</i> = -.02 (-.74, .71)	-5.27	<i>d</i> = -.78 (-1.62, .06)
<i>BDI-II</i>	TRAD vs. WL	-9.07*	<i>d</i> = -.89 (-1.64, -.14)	–	–	–	–
	EMMA vs. WL	-7.53*	<i>d</i> = -.78 (-1.52, -.04)	–	–	–	–
	EMMA vs. TRAD	1.53	<i>d</i> = .20 (-.52, .91)	-1.19	<i>d</i> = -.12 (-.84, .61)	-4.75	<i>d</i> = -.58 (-1.41, .24)
<i>PTGI</i>	TRAD vs. WL	12.43	<i>d</i> = .47 (-.26, 1.19)	–	–	–	–
	EMMA vs. WL	9.13	<i>d</i> = .34 (-.37, 1.06)	–	–	–	–
	EMMA vs. TRAD	-3.30	<i>d</i> = -.11 (-.82, .61)	-1.55	<i>d</i> = -.12 (-.85, .60)	2.24	<i>d</i> = .11 (-.70, .92)
<i>Impairment</i>	TRAD vs. WL	-2.06***	<i>d</i> = -.96 (-1.72, -.20)	–	–	–	–
	EMMA vs. WL	-1.43**	<i>d</i> = -.81 (-1.56, -.07)	–	–	–	–
	EMMA vs. TRAD	.62	<i>d</i> = .41 (-.32, 1.13)	-.30	<i>d</i> = -.23 (-.96, .51)	.51	<i>d</i> = -.41 (-1.23, .41)
<i>Severity</i>	TRAD vs. WL	-1.64**	<i>d</i> = -1.74 (-2.58, -.90)	–	–	–	–
	EMMA vs. WL	-2.25***	<i>d</i> = -1.94 (-2.81, -1.07)	–	–	–	–
	EMMA vs. TRAD	-.62	<i>d</i> = -.50 (-1.22, .23)	.072	<i>d</i> = .00 (-.73, .73)	.53	<i>d</i> = -.14 (-.96, .67)

Note: *d* = Cohen's *d*. Values between () represents the 95% Confidence Interval. Mean diff.= Mean differences. * $p < .05$, ** $p < .01$, *** $p < .001$. EMMA = EMMA Intervention group; TRAD= Traditional Intervention group; WL = Waiting list control group; ISL= Inventory of Stress and Loss; ICG= Inventory of Complicated Grief; BDI-II= Beck Depression Inventory II; PTGI= Posttraumatic Growth Inventory; Impairment= Maladjustment Scale; Severity= Clinician Severity Scale.

Table 4. Clinically Meaningful Improvement on the ISL/ICG and BDI at Post-treatment, 6 and 12 month follow up.

Measures	Post-treatment		6-month FW		12-month FW	
	EMMA	TRAD	EMMA	TRAD	EMMA	TRAD
ISL/ICG						
Recovered	10 (71.4%)	10 (66.7%)	12 (85.7%)	11 (84.6%)	13 (100%)	7 (70%)
Improved	0 (0%)	0(0%)	0 (0%)	0(0%)	0 (0%)	0 (0%)
No change	4 (28.6%)	5 (33.3%)	2 (14.3%)	2 (15.4%)	0 (0%)	3 (30%)
Impaired	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
BDI-II						
Recovered	4 (26.7%)	6 (40%)	8 (53.3%)	6 (42.9%)	10 (71.4%)	3 (30%)
Improved	1 (6.7%)	0 (0%)	1 (6.7%)	1 (7.1%)	0 (0%)	0 (0%)
No change	10 (66.7%)	9 (60%)	6 (40%)	7 (50%)	4 (28.6%)	7 (70%)
Impaired	0 (0%)	0(0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)

Table 5. Direct Scores from primary and secondary measures at pre, post, 6 and month follow-up and between-group comparisons at post-treatment.

	CG participants	PRE	POST	6-month FW	12-month FW	POST comparisons
<i>SL/ICG</i>						EMMA vs. TRAD vs WL $p = .304$
<i>EMMA</i>	P1	18	14	14	12	
<i>TRAD</i>	P2	22	18	–	–	
	P3	60	25	26	–	
<i>WL</i>	P4	28	34	–	–	
	P5	27	17	–	–	
	P6	14	54	–	–	
<i>BDI-II</i>						
<i>EMMA</i>	P1	11	8	7	4	$p = .456$
<i>TRAD</i>	P2	25	21	–	–	
	P3	35	12	5	–	
<i>WL</i>	P4	28	31	–	–	
	P5	2	8	–	–	
	P6	36	29	–	–	
<i>PTGI</i>						
<i>EMMA</i>	P1	13	53	44	52	$p = .304$
<i>TRAD</i>	P2	5	22	–	–	
	P3	39	75	69	–	
<i>WL</i>	P4	68	74	–	–	
	P5	68	77	–	–	
	P6	63	67	–	–	
<i>Impairment</i>						
<i>EMMA</i>	P1	5	2	2	2	$p = .753$
<i>TRAD</i>	P2	5	3	–	–	
	P3	6	3	2	–	
<i>WL</i>	P4	8	8	–	–	
	P5	5	1	–	–	
	P6	4	8	–	–	
<i>Severity</i>						
<i>EMMA</i>	P1	5	1	2	1	$p = .245$
<i>TRAD</i>	P2	5	2	–	–	
	P3	7	4	3	–	
<i>WL</i>	P4	6	5	–	–	
	P5	3	4	–	–	
	P6	6	5	–	–	

EMMA = EMMA Intervention group; TRAD= Traditional Intervention group; WL = Waiting list control group. CG. Complicated Grief. FW= Follow-up. POST Comparisons: Kruskal–Wallis test.




Figure 1. Flow diagram of the study participants, point of random assignments, and dropouts at each stage of the study




Figure 2. Scores for primary outcome measures at pre-, post-treatment and follow-ups. TRAD = Traditional Intervention group. EMMA = EMMA Intervention group. WL = Waiting list control group.

Accepted Article